

## Supplementary Online Content

Yaghi S, Boehme AK, Dibu J, et al. Treatment and outcome of thrombolysis-related hemorrhage: a multicenter retrospective study. Published online October 26, 2015. *JAMA Neurol*. doi:10.1001/jamaneurol.2015.2371.

**eTable.** Comparisons Between Those With and Without Follow-up Examinations

This supplementary material has been provided by the authors to give readers additional information about their work.

© 2015 American Medical Association. All rights reserved.

**eTable.** Comparisons Between Those With and Without Follow-up Examinations

	Without Follow up exam (N=46)	With Follow up exam (N=82)	p-value
Age in years (median, range)	76 (42-99)	73 (23-94)	0.0917
Gender (% male)	56.5%	45.1%	0.2158
<b><i>Ethnicity</i></b>			0.9502
Caucasian	63%	63.4%	
African American	30.4%	29.3%	
Hispanic	4.4%	6.1%	
Other	2.2%	1.2%	
NIHSS (median)	15 (0-33)	14 (1-32)	0.6836
Neuro-Interventional Procedure	13%	14.6%	0.8038
Hematoma Volume in cc (median, range)	23 (1-245)	25 (1-175)	0.5251
Time from rtPA to sICH diagnosis in hours (median, range)	442 (59-1980)	486 (30-2572)	0.3101
Time to treatment of sICH in hours (median, range)	83 (29-628)	112 (12-419)	0.5363
<b><i>Type of treatment</i></b>			
Any treatment	30%	52%	0.0404
Fresh Frozen Plasma	19.6%	20.7%	0.8749
Vitamin K	4.4%	3.9%	0.2988
Cryoprecipitate	28.3%	32.9%	0.3665
Platelets	28.3%	29.3%	0.1601

© 2015 American Medical Association. All rights reserved.

Amicar	0	3.7%	0.2594
Prothrombin Complex Concentrate	0	3.7%	0.2594
Factor VIIa	0	3.7%	0.2594
Surgical treatment	19.5%	12.2%	0.2791
Code Status Change in first 24 hours	43.5%	20.7%	0.0065
Mortality	69.6%	42.7%	0.0035

© 2015 American Medical Association. All rights reserved.