

Table S1. Baseline prevalence and risk of new onset of pain by location

	Prevalence ^a , n=2403			Risk ^b , n=374		
	Total	traumatic	Non-traumatic	Total	traumatic	Non-traumatic
Overall	634 (27.4)	129 (5.8)	509 (22.3)	82 (22.7)	11 (3.1)	74 (20.6)
Location						
Neck	56 (2.4)	5 (0.2)	50 (2.1)	9 (2.5)	2 (0.5)	6 (1.6)
Upper limbs	224 (9.5)	41 (1.8)	169 (7.2)	23 (6.3)	3 (0.8)	18 (4.9)
Chest	14 (0.6)	1 (0.04)	13 (0.6)	3 (0.8)	0 (0)	3 (0.8)
Upper back	34 (1.4)	2 (0.1)	30 (1.3)	1 (0.3)	0 (0)	1 (0.3)
Lower back	202 (8.5)	16 (0.7)	177 (7.5)	26 (7.0)	1 (0.3)	23 (6.2)
Buttocks	19 (0.8)	2 (0.1)	17 (0.7)	4 (1.1)	0 (0)	4 (1.1)
Lower limbs	360 (15.4)	77 (3.3)	257 (11.1)	44 (11.9)	5 (1.4)	38 (10.3)

Values are numbers (percentages). Sample sizes (denominators) of number counts vary due to missing values.

^a No duplication by student (1 record for each student).

^b Point prevalence (new onset) of pain at 1-year follow-up, among those who had no pain and in 7th or 10th grade at baseline.