

Sylt-Rømø Bight					North Sea				
trophic group	species	d13C (‰)	d15N (‰)	n	trophic group	species	d13C (‰)	d15N (‰)	n
planktivorous/ piscivorous	<i>Ammodytes tobianus</i>	-19.0 ± 1.2	16.0 ± 0.9	38	planktivorous/ piscivorous	<i>Hyperoplus lanceolatus</i>	-19.8 ± 0.9	16.2 ± 0.1	3
	<i>Belone belone</i>	-17.0 ± 1.1	16.5 ± 0.6	7		<i>Sprattus sprattus</i>	-21.0 ± 2.2	15.2 ± 0.1	2
	<i>Clupea harengus</i>	-18.8 ± 0.4	16.0 ± 0.7	54	Benthivorous/ piscivorous	<i>Ciliata mustela</i>	-18.2 ± 0.4	17.6 ± 0.6	2
	<i>Hyperoplus lanceolatus</i>	-18.4 ± 1.6	16.0 ± 1.3	14		<i>Gadus morhua</i>	-18.4 ± 0.5	17.7 ± 0.3	3
	<i>Sprattus sprattus</i>	-18.3 ± 1.9	15.8 ± 1.1	28		<i>Limanda limanda</i>	-20.0 ± 1.0	15.6 ± 0.2	6
Benthivorous/ piscivorous	<i>Gasterosteus aculeatus</i>	-18.9 ± 1.4	16.9 ± 0.6	9	<i>Merlangius merlangus</i>	-18.4 ± 1.3	16.9 ± 0.4	12	
	<i>Limanda limanda</i>	-16.9 ± 0.6	16.6 ± 1.0	18	<i>Agonus cataphractus</i>	-16.8 ± 0.1	17.4 ± 0.2	3	
	<i>Merlangius merlangus</i>	-17.1 ± 1.9	16.8 ± 1.3	36	<i>Crangon crangon</i>	-17.3 ± 2.2	16.2 ± 0.7	6	
	<i>Myoxocephalus scorpius</i>	-15.6 ± 0.6	17.1 ± 1.0	17	Strictly benthivorous	<i>Pomatoschistus microps</i>	-19.1	15.0	1
	<i>Platichthys flesus</i>	-16.2 ± 1.3	16.2 ± 1.7	8		<i>Pomatoschistus minutus</i>	-17.9 ± 0.2	16.3 ± 0.1	5
	<i>Syngnathus rostellatus</i>	-17.6 ± 1.0	16.8 ± 0.6	30		<i>Pleuronectes platessa</i>	-19.9 ± 0.9	14.7 ± 0.7	6
<i>Agonus cataphractus</i>	-15.9 ± 1.6	17.2 ± 0.9	19	<i>Solea solea</i>		-18.5 ± 0.7	16.4 ± 0.8	3	
Strictly benthivorous	<i>Crangon crangon</i>	-16.1 ± 1.2	16.1 ± 1.3	9	<i>Zoarces viviparus</i>	-17.8 ± 0.6	16.8 ± 0.5	9	
	<i>Pholis gunnellus</i>	-16.6 ± 2.8	17.0 ± 1.0	9					
	<i>Pomatoschistus microps</i>	-14.1 ± 1.6	17.0 ± 1.0	21					
	<i>Pomatoschistus minutus</i>	-16.9 ± 1.0	16.5 ± 0.4	42					
	<i>Pleuronectes platessa</i>	-15.7 ± 1.8	15.8 ± 0.5	61					
	<i>Zoarces viviparus</i>	-15.6 ± 1.6	16.7 ± 0.6	16					
	<i>loligo sp.</i>	-19.0 ± 0.8	14.0 ± 1.0	15					
<i>Osmerus eperlanus</i>	-16.5 ± 0.9	17.4 ± 0.7	20						