

Supplementary Table 2. Baseline characteristics by age

	12-21			22-35		
	Treatment (n= 26)	Placebo (n= 12)	Total 12-21 (n=38)	Treatment (n= 12)	Placebo (n= 8)	Total 22-35 (n=20)
Age in years	15.6 (2.72)	15.3 (2.23)	15.5 (2.54)	27.7 (4.48)	28.1 (3.87)	27.9 (4.15)
Men	18 (69.2%)	6 (50.0%)	24 (63.2%)	6 (50.0)	5 (62.5)	11 (55.0)
Ethnic origin						
White	21 (80.8%)	9 (75.0%)	30 (78.9%)	11 (91.7%)	8	19 (95.0%)
Non-white	5 (19.2%)	3 (25.0%)	8 (21.1%)	1 (8.3%)	0	1 (5.0%)
Body-mass index	22.489 (3.3942)	23.241 (2.4466)	22.726 (3.1128)	23.603 (3.2800)	26.047 (3.9608)	24.581 (3.6765)
Days since diagnosis	67.9 (21.31)	83.0 (17.33)	72.7 (21.14)	71.3 (20.97)	66.8 (15.22)	69.5 (18.58)
Baseline 2-hour C-peptide AUC (pmol/ml)	0.97 (0.390)	0.98 (0.565)	0.97 (0.445)	0.62 (0.161)	0.86 (0.415)	0.71 (0.306)
Baseline Peak C-peptide (pmol/ml)	1.24 (0.461)	1.22 (0.620)	1.23 (0.508)	0.91 (0.204)	1.13 (0.441)	1.00 (0.330)
Baseline HbA1c (%)	6.81 (0.966)	6.73 (1.405)	6.78 (1.104)	6.62 (1.804)	6.88 (0.997)	6.72 (1.506)
Baseline insulin use (U/kg per day)	0.37 (0.246)	0.47 (0.273)	0.41 (0.256)	0.26 (0.099)	0.33 (0.157)	0.29 (0.127)
Days of insulin use prior to randomization	68.8 (19.2)	83 (17.3)	73.3 (19.6)	71.3 (20.9)	66.8 (15.2)	69.5 (18.5)
GAD65 (% positive)	84.62%	91.67%	86.84%	75%	85.71%	78.95%
IA.2ic (% positive)	65.38%	50%	60.53%	58.33%	57.14%	57.89%
ZnT8 (% positive)	64%	50%	59.46%	66.67%	42.86%	57.89%
ICA (% positive)	61.90%	33.33%	51.52%	70%	100%	80%