

Supplementary Information

Chitosan oligosaccharide induces resistance to *Tobacco*

mosaic virus in Arabidopsis via salicylic acid-mediated signal pathway

Xiaochen Jia^{a, b, +}, Qingshan Meng^{a, c, +}, Haihong Zeng^{a, d}, Wenxia Wang^a, Heng Yin^{a, *}

^a Liaoning Provincial Key Laboratory of Carbohydrates, Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, China

^b University of Chinese Academy of Sciences, Beijing, 100049, China

^c College of Life Science, Dalian University of Technology, Dalian 116023, China

^d College of Food Science and Engineering, Dalian Ocean University, Dalian 116023, China

⁺ These authors contributed equally to this work

*Corresponding author at: Liaoning Provincial Key Laboratory of Carbohydrates, Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, China.

E-mail address: yinheng@dicp.ac.cn (H. Yin).




Figure S1. The full-length western-blot in Fig.1 and Fig.3. A for Fig.1A; B for Fig.1B; C for Fig.3A, These were the representative experiment and were independently repeated three times using different samples from independent treatment. The same amount of protein was loaded on SDS-PAGE gels, detected by anti-plant actin antibody, which served as the protein loading control. The detection of TMV-CP and loading control were carried out under the same experimental conditions.