

# Northcott et al. Supplementary Table 6

**A**

	Hazard ratio	.95 lower CI	.95 upper CI	logrank p-value
age18	1.05	0.51	2.167	0.900
Gender	1.33	0.85	2.064	0.209
Mstage.dM1	0.93	0.44	1.948	0.828
Mstage.dM2	1.18	0.47	2.937	0.726
Mstage.dM3	2.23	1.32	3.772	0.002
Resection	0.78	0.52	1.180	0.238
Histology.dDMB	0.82	0.30	2.242	0.686
Histology.dLGA	4.92	2.89	8.375	<0.001
WNT	0.52	0.26	1.042	0.061
SHH	0.57	0.34	0.956	0.031
C	3.97	2.55	6.194	<0.001
D	0.81	0.50	1.328	0.406

## Univariate prognostic value - PFS

**B**

	coef	exp(coef)	se(coef)	z	p
age18>18	0.284	1.328	0.383	0.740	0.459
GenderM	0.319	1.375	0.227	1.402	0.161
Resection1	-0.064	0.938	0.222	-0.287	0.774
HistologyDMB	-0.160	0.852	0.524	-0.305	0.761
HistologyLGA	1.492	4.448	0.297	5.028	0.000
MstageM1	-0.082	0.921	0.382	-0.215	0.830
MstageM2	-0.124	0.883	0.479	-0.259	0.795
MstageM3	0.500	1.649	0.291	1.718	0.086

## Clinical multivariate Cox PH model for PFS

**C**

	coef	exp(coef)	se(coef)	z	p
age18>18	0.731	2.078	0.446	1.639	0.101
GenderM	0.226	1.254	0.238	0.952	0.341
Resection1	-0.044	0.957	0.229	-0.191	0.848
HistologyDMB	0.270	1.311	0.538	0.503	0.615
HistologyLGA	1.330	3.780	0.299	4.453	0.000
MstageM1	-0.296	0.744	0.409	-0.723	0.469
MstageM2	-0.250	0.779	0.481	-0.519	0.604
MstageM3	0.262	1.299	0.295	0.886	0.376
SHH1	-0.072	0.930	0.485	-0.149	0.882
WNT1	-0.477	0.621	0.646	-0.738	0.460
C1	1.362	3.904	0.504	2.704	0.007
D1	0.270	1.310	0.488	0.553	0.580

## clinical/molecular multivariate Cox PH model for PFS