Compare Nursing Homes

General Information Inspections and Complaints Staffing Quality Measures Penalties		
	Nursing Home D	Nursing Home E
Overall Rating	Average	Average
Health Inspection Rating	Average	Below Average
Staffing Rating	Much Below Average	Above Average
Quality Measure Rating	Much Above Average	Below Average
LONG-STAY RESIDENTS Current data collection period		
Percent of long-stay residents experiencing one or more falls with major injury. Lower percentages are better.	0.0%	6.0%
Percent of long-stay residents with a urinary tract infection. Lower percentages are better.	3.4%	12.2%
Percent of long-stay residents who self-report moderate to severe pain. Lower percentages are better.	2.4%	4.6%
Percent of long-stay high-risk residents with pressure ulcers. Lower percentages are better.	4.9%	8.3%
Percent of long-stay low-risk residents who lose control of their bowels or bladder. Lower percentages are better.	23.1%	39.0%
Percent of long-stay residents who have/had a catheter inserted and left in their bladder. Lower percentages are better.	3.0%	4.9%
Percent of long-stay residents who were physically restrained. Lower percentages are better.	1.0%	2.1%
Percent of long-stay residents whose need for help with daily activities has increased. Lower percentages are better.	9.1%	4.9%
Percent of long-stay residents who lose too much weight. Lower percentages are better.	6.1%	11.8%
Percent of long-stay residents who have depressive symptoms. Lower percentages are better.	6.8%	1.1%
Percent of long-stay residents assessed and given, appropriately, the seasonal influenza vaccine. <i>Higher</i> percentages are better.	78.8%	95.8%
Percent of long-stay residents assessed and given, appropriately, the pneumococcal vaccine. <i>Higher</i> percentages are better.	91.6%	86.3%
Percent of long-stay residents who received an antipsychotic medication. Lower percentages are better.	13.4%	28.2%