

Nursing Home Profile

General Information

Inspections and Complaints

Staffing

Quality Measures

Penalties

Nursing Home D

Overall Rating



Average

Quality Measure



Much Above Average

Nursing Home D

State Average

National Average

LONG-STAY RESIDENTS

Current data collection period

	Nursing Home D	State Average	National Average
Percent of long-stay residents experiencing one or more falls with major injury. <i>Lower percentages are better.</i>	0.0%	3.4%	3.4%
Percent of long-stay residents with a urinary tract infection. <i>Lower percentages are better.</i>	3.4%	7.1%	7.6%
Percent of long-stay residents who self-report moderate to severe pain. <i>Lower percentages are better.</i>	2.4%	9.2%	11.6%
Percent of long-stay high-risk residents with pressure ulcers. <i>Lower percentages are better.</i>	4.9%	7.7%	6.8%
Percent of long-stay low-risk residents who lose control of their bowels or bladder. <i>Lower percentages are better.</i>	23.1%	34.4%	42.3%
Percent of long-stay residents who have/had a catheter inserted and left in their bladder. <i>Lower percentages are better.</i>	3.0%	4.7%	4.2%
Percent of long-stay residents who were physically restrained. <i>Lower percentages are better.</i>	1.0%	2.6%	2.2%
Percent of long-stay residents whose need for help with daily activities has increased. <i>Lower percentages are better.</i>	9.1%	15.1%	16.4%
Percent of long-stay residents who lose too much weight. <i>Lower percentages are better.</i>	6.1%	8.1%	7.3%
Percent of long-stay residents who have depressive symptoms. <i>Lower percentages are better.</i>	6.8%	7.7%	7.4%
Percent of long-stay residents assessed and given, appropriately, the seasonal influenza vaccine. <i>Higher percentages are better.</i>	78.8%	92.6%	93.4%
Percent of long-stay residents assessed and given, appropriately, the pneumococcal vaccine. <i>Higher percentages are better.</i>	91.6%	92.9%	94.3%
Percent of long-stay residents who received an antipsychotic medication. <i>Lower percentages are better.</i>	13.4%	26.0%	23.8%