

Table S1. ICD-10-AM codes used in the identification of sleep apnea

ICD-10-AM code	Description	Number of women (% of all cases in study)
G47.30	Sleep apnoea, unspecified	170 (32.8%)
G47.31	Central sleep apnoea syndrome (Central sleep hypopnoea syndrome)	6 (1.2%)
G47.32	Obstructive sleep apnoea syndrome (Obstructive sleep hypopnoea syndrome)	336 (64.7%)
G47.33	Sleep hypoventilation syndrome	1 (0.2%)
G47.39	Other sleep apnoea	6 (1.2%)
Total		519 (100.0%)

Note: ICD-10-AM refers to the Australian Modification of the ICD-10. These codes differ from those in the ICD-10.

Table S2. ICD-10-AM codes used in the identification of obesity

ICD-10-AM code	Description	Number of records (% of all records of obesity)
E65	Localised adiposity	156 (4.1%)
E66.0	Obesity due to excess calories	12 (0.3%)
E66.1	Drug-induced obesity	1 (0.0%)
E66.2	Extreme obesity with alveolar hypoventilation	None
E66.8	Other obesity (Morbid obesity)	1,282 (33.4%)
E66.9	Obesity, unspecified (Simple obesity Not Otherwise Specified)	2,388 (62.2%)
Total		3,839 (100.0%)

Note: There are 3 more obesity records than women with obesity (N = 3,836) because 2 women had both E65 and E66.9, and 1 woman had both E66.1 and E66.9.