

Common *UCP2* variants contribute to serum urate concentrations and the risk of hyperuricemia

Luyu Yang^{1¶}, Zheng Dong^{1¶}, Jingru Zhou¹, Yanyun Ma¹, Weilin Pu¹, Dongbao Zhao², Hongjun He³,
Hengdong Ji⁴, Yajun Yang^{1,5}, Xiaofeng Wang^{1,5}, Xia Xu², Yafei Pang², Hejian Zou^{6,7}, Li Jin^{1,5},
Chengde Yang^{8*}, Jiucun Wang^{1,5,7*}

¹State Key Laboratory of Genetic Engineering and Ministry of Education Key Laboratory of Contemporary Anthropology, Collaborative Innovation Center for Genetics and Development, School of Life Sciences, Fudan University, Shanghai, China.

²Division of Rheumatology and Immunology, Changhai Hospital, Shanghai, China.

³Division of Rheumatology, Taixing People's Hospital, Jiangsu Province, China.

⁴Division of Rheumatology, Taizhou People's Hospital, Jiangsu Province, China.

⁵Fudan-Taizhou Institute of Health Sciences, Taizhou, Jiangsu Province, China.

⁶Division of Rheumatology, Huashan Hospital, Fudan University, Shanghai, China.

⁷Institute of Rheumatology, Immunology and Allergy, Fudan University, Shanghai, China.

⁸Division of Rheumatology, Ruijin Hospital, Shanghai Jiaotong University School of Medicine, Shanghai, China.

*Corresponding author

E-mail: yangchengde@sina.com (Chengde Yang), jcwang@fudan.edu.cn (Jiucun Wang)

¶These authors equally contributed to this study.

S.Table 1. Characteristics of hyperuricemia and healthy control*

	Control	Hyperuricemia
Number	2945	1387
Male/Female	2077/868	1060/327
Age (year)	69.00±9.25	67.91±11.93
Height (cm)	161.37±7.52	162.63±7.54
Weight (kg)	63.71±9.96	68.56±11.12
BMI	24.43±3.24	25.88±3.52
Smoking	87.05%	72.33%
Serum Urate (umol/L)	300.73±63.13	482.30±61.14

*All data shown in mean±SD