Oligomerization of the Microtubule Associated Protein Tau is Mediated by its N-Terminal Sequences: Implications for Normal and Pathological Tau Action

H. Eric Feinstein^{1,2,*}, Sarah J. Benbow^{1,2,*}, Nichole E. LaPointe^{1,*}, Nirav Patel^{3,*}, Srinivasan Ramachandran^{3,*}, Thanh D. Do^{4,6*}, Michelle R. Gaylord^{1,2,7}, Noelle E. Huskey^{1,2,8}, Nicolette Dressler⁵, Megan Korff⁵, Brady Quon⁵, Kristi Lazar Cantrell⁵, Michael T. Bowers⁴, Ratnesh Lal^{3,**}, Stuart C. Feinstein^{1,2,**}

Corresponding Author:
Dr. Stuart Feinstein
Neuroscience Research Institute
University of California
Santa Barbara CA 93106:
Tel: (805) 893-2659

Fax: (805) 893-2005 feinstei@lifesci.ucsb.edu

SUPPLEMENTAL MATERIALS

¹Neuroscience Research Institute and Department of Molecular, Cellular and Developmental Biology, University of California, Santa Barbara, CA 93106

²Department of Molecular, Cellular and Developmental Biology, University of California, Santa Barbara, CA 93106

³Department of Bioengineering, Department of Mechanical Engineering and Materials Science Graduate Program, University of California, San Diego, CA

⁴Department of Chemistry and Biochemistry, University of California, Santa Barbara, CA 93106

⁵Department of Chemistry, Westmount College, Santa Barbara, CA 93108


⁶ Current Address: Department of Chemistry and the Beckman Institute, University of Illinois at Urbana-Champaign, Urbana, 61801.

⁷Current Address: Genomics Institute of the Novartis Research Foundation 10675 John Jay Hopkins Drive San Diego, 92121

⁸Current Address: Department of Dermatology and Program in Epithelial Biology, Stanford University School of Medicine, Stanford, California, USA.

^{*}Co-first authors

^{**}Co-senior authors



Supplementary Figure S1. SDS gel analysis of all recombinant full-length and deletion tau constructs used in this work. 1 μ g of each sample (except tau⁴⁵⁻²³⁰) was fractionated on a 12% denaturing polyacrylamide gel to assess its purity. 5 μ g was used for tau⁴⁵⁻²³⁰.