

Table S1. Primers used for qRT-PCR

GENE	FORWARD	REVERSE
GAPDH	AGGGGTCTACATGGCAACTG	GGGTTTTCCAGAACCACAGA
CD31	GCTGAAGGCTGAAAGACAGG	GGCTGGTCTCAAACCTCCTGA
CD34	TTTGGGCCCTCTCCCATCC	GCCGGTCCCGTTTTCTGAG
VE-cadherin	CCGAGCCATGTGTCTTTACA	TGAGCTTGGTGTCTCCTTCC
GATA2	CAGACGAAGGCAACCATTTT	CAGAGGAGAAGAGGGTGCAG
TAL1 UTR	GTCCCAAATCCTTCCCAAGT	GCTGACAACCCAGGTCTTA
FLI1 UTR	AGCGAGCTGGTCTATCCAGA	GCGTTATTCTTTGCTGGAG
FLI1 coding	GAGGGGGTTGATCTTGTGG	GGATGCCTTCCCTATGTTCA
SOX1	AGAACCCCAAGATGCACAAC	GCCAGCGAGTACTTGTCTT
SOX2 UTR	TTGTTCGATCCCAACTTTCC	ACATGGATTCTCGGCAGACT
HOPX UTR	GGAGGAGACCCAGAAATGGT	AACAGAGATGGCCTTCATGG
GFAP	TCTCTCGGAGTATCTGGGAAGT	TTCCCTTCTGTCTGAGTCTCA
MAP2	GGGCCTTTTCTTTGAAATCTAGTTT	CAAATGTGGCTCTCTGAAGAACA
PAX6	CCAGCCAGACCTCCTCATAC	TCACTTCCGGGAAGTGAAC
N-cadherin	TGTGGGAATCCGACGAATGGATGA	TGGAGCCACTGCCTTCATAGTCAA
CD44	TCTCAAGGGCGTAACTCTGG	GCCAATTCTACCAGGCTTGA
CD73	TCTGCCTCCAAATCTGAACA	TCTAGCTGCCATTTGCACAC
LIN28B	GCCCCTTGGATATTCCAGTC	TGACTCAAGGCCTTTGGAAG
HMGA2	ACTTCAGCCCAGGGACAAC	TTGAGCTGCTTTAGAGGGAC