

SUPPLEMENTAL DIGITAL CONTENT 3

TABLE. Mean percent differences in FA values for healthy controls.

<i>Bundle</i>	<i>C vs E₁</i>	<i>C vs E₂</i>	<i>C vs E</i>	<i>E₁ vs E₂</i>	<i>E₁ (Intra)</i>	<i>E₂ (Intra)</i>	<i>E (Intra)</i>
<i>I</i>FOF	3.26 (0.70)	3.90 (1.11)	3.58 (0.98)	5.16 (1.18)	4.65 (1.03)	7.32 (1.47)	5.98 (1.84)
<i>I</i>LF	2.58 (1.68)	1.50 (0.36)	2.04 (1.33)	2.70 (0.82)	1.89 (0.28)	2.36 (0.31)	2.12 (0.38)
<i>A</i>F	3.58 (1.63)	1.95 (0.82)	2.76 (1.53)	4.28 (2.30)	0.80 (0.30)	1.17 (0.37)	0.98 (0.39)
<i>U</i>F	3.46 (2.47)	2.51 (1.66)	2.98 (2.16)	2.00 (1.30)	1.17 (1.06)	2.03 (1.17)	1.60 (1.20)
<i>C</i>B	1.16 (0.30)	1.41 (0.58)	1.29 (0.48)	0.64 (0.14)	1.00 (0.14)	0.33 (0.06)	0.66 (0.35)
<i>F</i>R	1.67 (0.44)	1.25 (0.68)	1.46 (0.61)	2.39 (0.17)	1.76 (1.60)	1.47 (1.54)	1.62 (1.58)

The standard deviations are given in parentheses. C, clustering; E₁, expert 1; E₂, expert 2; E, both experts (average of both). Intra-expert agreements quantify the difference between the repeated drawings. The overall difference between clustering and experts (C vs E) is comparable to the difference between experts (E₁ vs E₂).