Drugs tables

ADULTS

Category 1 (6 months) Regimen for New Cases: 2RHZE/4RH

Regimen	Pre-treatment weight			
	>70kg	55-70kg	38-54kg	21-37kg
Intensive phase (2 months):	5	4	3	2
Combined tablet of RHZE (150mg+75mg+400mg+275mg)				
Continuation phase (4months):	5	4	3	2
Combined tablet of RH (150mg + 75mg)				

Category 1 (8months) Regimen for New Cases: 2 RHZE/ 6EH

Regimen	Pre-treatment weight			
	>70kg	55-70kg	38-54kg	21-37kg
Intensive phase (2 months):	5	4	3	2
Combined tablet of RHZE (150mg+75mg+400mg+275mg)				
Continuation phase (6months):	3	3	2	1
Combined tablet of EH (400mg + 150mg)				

CHILDREN

Category 1 Regimen for New Cases: 2 RHZ+E/ 4EH (0-14years)

Regimen	Pi	Pre-treatment weight			
	15-20kg	8-14kg	<7kg		
Intensive phase (2 months):	3	2	1		
Combined tablet of RHZ (60mg+30mg+150mg)	3	2	1		
Ethambutol 100mg					
Continuation phase (4months):	3	2	1		
Combined tablet of RH (60mg + 30mg)					

The CAT 2 regimen remains 8 months: daily supervision

ADULTS

Category 2 Regimen for Relapses, Failure, RAD and Others: 2SRHZE/RHZE/5RHE

Regimen	Pre-treatment weight			
	>70kg	55-70kg	38-54kg	21-37kg
Intensive phase (3 months):	5	4	3	2
Combined tablet of RHZE (150mg+75mg+400mg+275mg)				
Add in the first 2 months daily:	1 gram	1 gram	0.75 gram	0.5 gram
Streptomycin				
Continuation phase (5months):	5	4	3	2
Combined tablet of RHE (150mg+75mg+275mg)				

- i. Streptomycin should NOT be given to pregnant women.
- ii. Patients >45 years should not be given more than 0.75g of streptomycin irrespective of weight