## Bisdemethoxycurcumin inhibits ovarian cancer via reducing oxidative stress mediated MMPs expression

Haifeng Pei<sup>1</sup>, Yi Yang<sup>1</sup>, Lin Cui<sup>2</sup>, Jiong Yang<sup>1</sup>, Xiuchuan Li<sup>1</sup>, Yongjian Yang<sup>1,\*</sup>, Haixia Duan<sup>3,\*</sup>

<sup>1</sup> Department of Cardiology, Chengdu Military General Hospital, Chengdu 610083, China;

<sup>2</sup> Department of Orthopedics, Chengdu Military General Hospital, Chengdu 610083, China;

<sup>3</sup> Department of Obstetrics and Gynecology, Affiliated Guangren Hospital, College of Medicine,

Xi'an Jiaotong University, Xi'an 710004, China.

Running title: BDMC and Ovarian Cancer

## \*Address correspondence to:

Haixia Duan, MD, PhD Affiliated Guangren Hospital, College of Medicine, Xi'an Jiaotong University 21 Jiefang Road Xi'an 710004, China E-mail: dhx20167@126.com Tel: +86-29-8742-0006 Fax: +86-29-8742-0008 Or Yongjian Yang, MD, PhD Chengdu Military General Hospital 270 Tianhui Road Chengdu 610083, China E-mail: yyj10001@126.com Tel: +86-28-8657-0211 Fax: +86-28-8357-2211

Supplemental Table 1. The influence of BDMC on the cycle of SKOV-3 cell. BDMC arrests the cells at G1 phase of the cell cycle. Serum-1640 SKOV-3 ( $1 \times 10^{6}$  cells/ml) was incubated in the absence or presence of different doses of BDMC for 6, 12 and 24h, respectively. Thereafter, the cells were washed, fixed, stained with propidium iodide, and analyzed for DNA content by flow cytometry. BDMC, Bisdemethoxycurcumin. \*\*p<0.01 vs. 0  $\mu$ M BDMC.

BDMC (concentration) (µM)	Action time ( h )	Proportion of cell cycle (%)+		
		G1	s	G2
_	0	58.1	35.2	6.73+
5	6	60.5**	32.2	7.29+
10	6	65.6**	22	7.94+
15	6	66.5**	26.7	6.72+
5	12	70.1**	21.6	8.27
10	12	71.7**	20.1	8.16+
15	12	75.4**	19.7	4.86+
5	24	76.7**	17.3	6.03+
10	24	78.5**	15.6	5.85+
15	24	79.7**	16.4	3.89 +

Table 1. The influence of BDMC on the cycle of SKOV-3 cell.+