

Supplemental Table 2. Descriptive statistics for trauma history, appraisals, and PTSS

	Study 1	Study 2	Study 3	Total sample
TIME 1 (within 4 weeks)	<i>M (SD) days</i>	<i>M (SD) days</i>	<i>M (SD) days</i>	<i>M (SD)days</i>
	17.5 (8.4)	2.5 (2.6)	1.2 (1.5)	7.4 (9.4)
	N (%)	N (%)	N (%)	N (%)
Presence of trauma history	222 (80.4)	170 (58.2)	88 (73.3)	480 (69.8)
Appraisals				
Felt really afraid	111 (40.2)	186 (63.7)	94 (78.3)	391 (56.8)
Thought might die	57 (20.7)	41 (14.0)	26 (21.7)	124 (18.0)
Presence of significant PTSS	64 (23.2)	68 (23.3)	-	132 (19.2)
TIME 2 (within 6 weeks)	<i>M (SD) days</i>	<i>M (SD) days</i>	<i>M (SD) days</i>	<i>M (SD) days</i>
	-	55.6 (20.5)	27.9 (10.9)	45.8 (22.1)
	N (%)	N (%)	N (%)	N (%)
Presence of significant PTSS	-	38 (13.0)	11 (9.2)	49 (7.1)
Coping	<i>M (SD)</i>	<i>M (SD)</i>	<i>M (SD)</i>	<i>M (SD)</i>
Escape Coping*	-	4.48 (2.00)	4.37 (1.68)	4.44 (1.89)
Control Coping**	-	3.45 (1.34)	3.50 (1.20)	3.47 (1.29)
TIME 3 (3-6 months)	<i>M (SD) days</i>	<i>M (SD) days</i>	<i>M (SD) days</i>	<i>M (SD)days</i>
	209.4 (78.1)	201.5 (19.5)	112.9 (20.4)	199.1 (64.3)
	N (%)	N (%)	N (%)	N (%)
Presence of significant PTSS	39 (14.1)	29 (9.9)	17 (14.2)	85 (12.4)

Notes. Escape coping has a range of 0-10. Control coping has a range of 0-5.