

ANNEX: SUPPLEMENTARY MATERIAL.

Table S1. Respondents' perceptions of cancer risk factors: results from principal component analysis (contributions to the 3 first axes) (France, 2010, n =3,359).

	Contributions after varimax rotation ^a		
	Axis 1	Axis 2	Axis 3
Behavioral risk factors			
Tobacco smoking	0.00	0.22	0.02
Alcohol drinking	0.00	0.24	0.00
Having sun exposure without protection	0.00	0.20	0.02
Lack of physical activity	0.04	0.12	0.01
Having tanning lamp sessions	0.00	0.15	0.06
Cannabis smoking	0.00	0.14	0.02
Environmental risk factors			
Eating chemically-treated food	0.01	0.06	0.22
Breathing a polluted air	0.00	0.11	0.18
Living near a nuclear power plant	0.00	0.00	0.53
Living near a mobile telephone relay station	0.01	0.01	0.38
Psychosocial risk factors			
Suffering from stress of the modern life	0.14	0.03	0.02
Being weakened by painful experiences	0.17	0.00	0.01
Having difficulties in expressing feelings and emotions	0.15	0.00	0.00
Bitterness due to personal/professional disappointments	0.17	0.00	0.01

^a A varimax rotation was performed to maximize the sum of the variances of the squared loadings on each axe.