

Supplemental information

Multiple CDK inhibitor Dinaciclib suppresses neuroblastoma growth via inhibiting CDK2 and CDK9 activity

Zhenghu Chen^{1,2}, Zhenyu Wang^{3,4}, Jonathan C. Pang², Yang Yu², Shayahati Bieerkehazhi^{3,5}, Jiaxiong Lu^{1,2}, Ting Hu^{2,6}, Yanling Zhao², Xin Xu², Hong Zhang³, Joanna S. Yi², Shangfeng Liu^{1,7*}, and Jianhua Yang^{2*}

¹Department of Ophthalmology, Shanghai Tenth People's Hospital, Tongji University School of Medicine, Shanghai 200072, P. R. China

²Texas Children's Cancer Center, Department of Pediatrics, Dan L. Duncan Cancer Center, Baylor College of Medicine, Houston, Texas 77030, USA

³Department of Pathology, University of Texas MD Anderson Cancer Center, Houston, Texas 77030, USA

⁴Department of Breast Surgery, The second hospital of Jilin University, Changchun, Jilin 130041, China


⁵College of Public Health, Xinjiang Medical University, Urumqi 830054, China

⁶Department of Neurosurgery, Xiangya Hospital, Central South University, Changsha 410008, China


⁷Department of Stomatology, Huashan Hospital, Fudan University, Shanghai 200040, China

*Correspondence to:


Shangfeng Liu, Department of Stomatology, Huashan Hospital, Fudan University, Shanghai 200040, China. Tel/fax: +86-21-6562662. Email: shangfengliufudan@163.com. Jianhua Yang, Texas Children's Cancer Center, Department of Pediatrics, Dan L. Duncan Cancer Center, Baylor College of Medicine, Houston, TX 77030. USA. Phone: 1-832-824-4572, Fax: 1-832-825-1206. Email: jianhuay@bcm.edu;




Supplemental Figure S1. Correlations of the expressions of CDK2 and CDK9 with the outcomes in NB patients. (a) Relapse-free survival probability for NB patients with high CDK2 expression (blue; n=77) and low CDK2 expression (red; n=25) (Seeger-102 data set). (b) Relapse-free survival probability for NB patients with high CDK9 expression (blue; n=9) and low CDK9 expression (red; n=93) (Seeger-102 data set).




Supplemental Figure S2. Correlations of the expressions of CDK1 with the outcomes in NB patients. (a) Overall survival probability for NB patients with high CDK1 expression (blue; n=31) and low CDK1 expression (red; n=57) (Versteeg-88 data set). (b) Relapse-free survival probability for NB patients with high CDK1 expression (blue; n=31) and low CDK1 expression (red; n=57) (Versteeg-88 data set).



Supplemental Figure S3. Correlations of the expressions of CDK5 with the outcomes in NB patients. (a) Overall survival probability for NB patients with high CDK5 expression (blue; n=11) and low CDK5 expression (red; n=77) (Versteeg-88 data set). (b) Relapse-free survival probability for NB patients with high CDK5 expression (blue; n=8) and low CDK5 expression (red; n=80) (Versteeg-88 data set).



Supplemental Figure S4. The cytotoxic effects of CDK1/5 inhibitor and CDK2/9 inhibitor on NB-19 and SK-N-AS cells. NB-19 (a) and SK-N-AS (b) cells were treated with increasing concentrations of either CDK1/5 dual inhibitor or CDK2/9 dual inhibitor for 48 hrs. Cell viability was then measured by the Cell Counting Kit-8 (CCK-8) assay. P-values <0.01 (**) or P <0.001 (***) (Student's t-test, two-tailed) were indicated.



Supplemental Figure S5. The comparison of the cytotoxic effects caused by CDK inhibitors on NB-19 and SK-N-AS cells. NB-19 (a) and SK-N-AS (b) cells were treated with increasing concentrations of CDK1/5 dual inhibitor, or CDK2/9 dual inhibitor, or their combinations, or Dinaciclub for 36 hrs. Cell viability was then measured by the Cell Counting Kit-8 (CCK-8) assay. P-values <0.001 (***) (Student's t-test, two-tailed) were indicated.