

Supplementary Data

SUPPLEMENTARY TABLE S1. DISTRIBUTION OF OUTCOMES IN THE TREATED GROUP ACCORDING TO TSH LEVELS DURING THE SECOND HALF OF PREGNANCY^a

<i>Maternal outcomes, n (%)</i>	<i>LT4 group with TSH at goal, n=50</i>	<i>LT4 group with TSH not at goal, n=12</i>	<i>p-Value unadjusted</i>
Pregnancy loss	0 (0)	0 (0)	0.99
Preterm delivery (<37 weeks)	2 (4)	2 (16.7)	0.17
Gestational diabetes	1 (2)	1 (8.3)	0.35
Gestational hypertension	5 (10)	2 (16.7)	0.61
Pre-eclampsia	2 (4)	0 (0)	0.99
Premature rupture of membranes	6 (12)	1 (8.3)	0.99
Intrauterine growth restriction	1 (2)	0 (0)	0.99
Placenta previa	0 (0)	0 (0)	0.99
Placenta abruption	0 (0)	0 (0)	0.99
<i>Neonatal outcomes, n (%)</i>			
NICU admission	1 (2)	1 (8.3)	0.36
Birth weight < 2500 g	0 (0)	1 (8.3)	0.19
5-minute Apgar ≤ 7	0 (0)	0 (0)	0.99
Congenital malformations	0 (0)	0 (0)	0.99
Neonatal death	0 (0)	0 (0)	0.99

^aOut of the 82 treated pregnant women, 72 had a TSH level checked in the second half of the pregnancy (>20th gestational week); 50 had TSH valued ≤ 3 mIU/L (at goal) at all checks, 12 had TSH values >3 mIU/L at all checks, and 10 had TSH values fluctuating in and out of goal (excluded from this analysis).