

Online Table 1: Biopsy Site and Organ Transplantation in symptomatic subjects

	USA			Rest of World		
	Overall (n=330)	Cardiology Sites (n=309)	Non-Cardiology Sites (n=21)	Overall (n=1541)	Cardiology Sites (n=290)	Non-Cardiology Sites (n=1251)
Tissue type (number of biopsies)	353	338	15	1356	251	1105
Abdominal fat pad/fat pad	55 (15.6%)	54 (16.0%)	1 (6.7%)	228 (16.8%)	31 (12.4%)	197 (17.8%)
Cardiac	226 (64.0%)	220 (65.1%)	6 (40.0%)	174 (12.8%)	151 (60.2%)	23 (2.1%)
GI (except rectum)	16 (4.5%)	14 (4.1%)	2 (13.3%)	96 (7.1%)	21 (8.4%)	75 (6.8%)
Nerve	12 (3.4%)	10 (3.0%)	2 (13.3%)	108 (8.0%)	13 (5.2%)	95 (8.6%)
Skin	2 (0.6%)	1 (0.3%)	1 (6.7%)	55 (4.1%)	2 (0.8%)	53 (4.8%)
Rectal	4 (1.1%)	4 (1.2%)	0	48 (3.5%)	23 (9.2%)	25 (2.3%)
Salivary gland	0	0	0	579 (42.7%)	1 (0.4%)	578 (52.3%)
Other*	39 (10.8%)	35 (10.4%)	3 (20.0%)	68 (4.85.0%)	9 (3.6%)	59 (5.3%)
Transplants						

Liver Transplant	11 (3.3%)	9 (2.9%)	2 (9.5%)	286 (18.6%)	51 (17.6%)	235 (17.918.8%)
Heart Transplant	11 (3.3%)	8 (2.6%)	3 (14.3%)	16 (1.0%)	13 (4.5%)	3 (0.2%)
*Other locations for biopsy include stomach, small bowel, bone marrow, eye, eye vitrectomy, liver, lung, lymph node, median nerve/carpal tunnel tissue, sigmoid, synovial and transynovial tissue, thyroid, and transverse carpal ligament.						