

Appendix Table 1 : Pregnancies (live births, late fetal losses after 20 weeks gestation and terminations) with an NTD in England, Wales, Scotland and Ireland 1991-2012

Year of birth	Prevalence of all pregnancies with an NTD (adj for register)	Prevalence of live births with an NTD (adj for register)	Number of population live births (1,000s)	Number of pregnancies with an NTD	Proportion of pregnancies with an NTD resulting in a termination (%)	Number of terminations with an NTD	Number of live births with an NTD
	A	B			C		
1991	1.61	0.28	792	1274	81	1029	221
1992	1.54	0.26	781	1206	81	981	200
1993	1.67	0.17	762	1274	85	1083	126
1994	1.24	0.13	750	928	86	798	95
1995	1.39	0.25	732	1020	77	789	186
1996	1.41	0.14	733	1030	88	907	105
1997	1.50	0.21	727	1088	80	866	153
1998	1.31	0.13	717	936	85	797	96
1999	1.28	0.17	700	898	81	732	119
2000	1.32	0.17	679	898	84	751	118
2001	1.32	0.16	669	884	83	738	108
2002	1.37	0.22	669	915	78	714	144
2003	1.34	0.17	696	931	82	760	120
2004	1.27	0.18	716	910	81	739	132
2005	1.26	0.16	716	905	85	766	115
2006	1.26	0.17	742	931	83	774	129
2007	1.13	0.16	772	875	83	722	125
2008	1.09	0.21	794	863	78	676	166
2009	1.29	0.21	790	1019	80	814	166
2010	1.27	0.25	807	1022	77	787	201
2011	1.28	0.18	808	1034	82	851	142
2012	1.24	0.21	813	1012	76	772	172