

Supplementary Table 1. Incubation periods estimated using non-parametric and parametric methods according to the disease generation, a link to superspreaders, days of illness in source cases.

	Non-parametric Estimation	Parametric Estimation		
	Median days (IQR)	Median days (95% CI)	5 Percentile (95% CI)	95 Percentile (95% CI)
Generation				
1st generation (n=28)	5 (4-7)	6.2 (5.4-7.0)	3.2 (2.6-14.0)	10.6 (8.3-12.7)
2nd generation (n=113)	7 (5-10.5)	7.7 (7.0-8.4)	2.9 (2.50-3.4)	16.0 (14.3-17.7)
3rd generation (n=21)	8 (5-10)	7.95 (6.3-10.0)	2.8 (1.82-4.8)	17.2 (13.3-21.2)
Link to superspreaders				
SSE (n=149)	6 (4.5-9.5)	7.3 (6.7-7.9)	2.9 (1.69-8.7)	14.9 (13.4-16.4)
Non-SSE (n=13)	10 (8-13)	9.5 (7.1-12.0)	3.5 (1.69-8.7)	19.9 (15.0-23.0)
Days of illness in source cases				
≤3 days of illness (n=30)	9 (5.5-12)	8.9 (7.2-10.7)	3.7 (2.60-5.3)	21.6 (16.9-26.5)
4-6 days of illness (n=112)	6.75 (4.5-9.5)	7.3 (6.7-7.9)	3.0 (2.5-3.5)	14.5 (12.9-16.0)
≥7days of illness (n=20)	5 (3.26-8.5)	6.0 (4.8-7.3)	2.4 (1.9-3.4)	12.1 (9.1-14.6)

IQR, interquartile range; CI, confidence interval; SSE, superspreading events.