

## SUPPLEMENTARY INFORMATION

### Classification of Colon Cancer Patients Based on the Methylation Patterns of Promoters

Wonyoung Choi<sup>1†</sup>, Jungwoo Lee<sup>1†</sup>, Jin-Young Lee<sup>2</sup>, Sun-Min Lee<sup>2</sup>,  
Da-Won Kim<sup>1</sup>, Young-Joon Kim<sup>1,2\*</sup>

<sup>1</sup>Department of Integrated OMICS for Biomedical Science, Graduate School,  
Yonsei University, Seoul 03722, Korea

<sup>2</sup>Department of Biochemistry, College of Life Science & Biotechnology,  
Yonsei University, Seoul 03722, Korea

**Supplementary Table 1.** Clinical data of three patients groups in colon cancer

		Group 1 (n = 110)	Group 2 (n = 76)	Group 3 (n = 109)
Race	Asian	4	5	2
	Black or African American	32	10	14
	White	57	80	75
Age (yr)	≥ 60	73	80	50
	< 60	27	18	50
Gender	Male	50	56	55
	Female	50	43	45
Tumor stage	I	16	24	7
	II	35	46	39
	III	32	25	30
	IV	18	4	18
Tumor size	T1	2	3	3
	T2	15	24	7
	T3	64	66	77
	T4	18	7	13
Microsatellite instability	Yes	0.9	12	0.9
	No	23	24	32
Metastasis	Yes	16	4	18
	No	65	74	63
Vital status	Alive	85	84	86
	dead	14	15	14
Residual tumor	R0	71	55	69
	R1	2	1	0
	R2	2	0	3
New tumor event after initial treatment	Yes	5	0	4
	No	17	7	6
Anatomic neoplasm subdivision	Ascending colon	16	29	14

Cecum	29	42	11
Descending colon	5	3	6
Hepatic flexure	7	8	4
Sigmoid colon	32	4	46
Transverse colon	7	7	13

---