

Direct Synthesis of Alkenyl Boronic Esters from Unfunctionalized Alkenes: A Boryl-Heck Reaction

William B. Reid, Jesse J. Spillane, Sarah B. Krause and Donald A. Watson*

Department of Chemistry and Biochemistry, University of Delaware,
Newark, Delaware 19716, United States

Supporting Information

Index	Page
1. General Experimental Details:	S2
2. Instrumentation and Chromatography:	S2
3. Synthesis of Non-commercial Starting Materials and Reagents:	S3
4. General Procedure for the α -Olefin Boryl-Heck Reaction:	S5
5. General Procedure for the Styrene Boryl-Heck Reaction:	S11
6. Additional Optimization Data:	S16
7. Computational and Experimental Mechanistic Insight:	S18
8. Crystallographic Details:	S21
9. References:	S22
10. Spectral Data:	S23

1. General Experimental Details:


Diethyl ether, tetrahydrofuran and dichloromethane were dried on alumina according to published procedures.¹ Catechol was purchased from Alfa Aesar and recrystallized from hot toluene prior to use. Boron trichloride was purchased from Acros as a 1 M solution in hexanes and used as received. N,N-Dicyclohexylmethylamine was purchased from TCI, distilled from calcium hydride (80 °C, 150 mtorr) and stored at rt on the bench in a nitrogen-filled Strauss flask. Trifluorotoluene was purchased from Sigma Aldrich in an anhydrous septum sealed bottle, transferred to a Straus flask by cannula transfer and sparged with nitrogen for 15 minutes. Lithium trifluoromethanesulfonate was purchased from Oakwood and stored in a nitrogen-filled glovebox or in a desiccator under air. Bis(trimethylsilylmethyl)-(cycloocta-1,5-diene) palladium(II) [(COD)Pd(CH₂SiMe₃)₂] was prepared according to a published procedure.² Tris(dibenzylideneacetone)dipalladium(0) [Pd₂(dba)₃] was purchased from Aldrich or Strem and used as received. Bis(3,5-di-*tert*-butylphenyl)(*tert*-butyl)phosphine (**L1**) was prepared according to a published procedure.³ Substrates ethyl 4-vinylbenzoate,⁴ 5-vinylbenzo-1,3-dioxole⁵ and 1-(toluene-4-sulfonyl)-2-vinyl-1*H*-indole⁶ were prepared according to published literature procedures. All other substrates and reagents were purchased in highest analytical purity from commercial suppliers and used as received. Vials used in the glovebox were dried in a gravity oven at 140 °C for a minimum of 12 h, transferred into the glovebox hot, and then stored at rt in the glovebox prior to use. All other glassware was flame-dried under vacuum prior to use. “Double manifold” refers to a standard Schlenk-line gas manifold equipped with nitrogen and vacuum (ca. 100 mtorr). All optimization reactions (0.25 mmol) were run in a nitrogen-filled glovebox and heated using an aluminum block on a magnetic stir plate. All yields in optimization reactions were determined using ¹H NMR using 1,3,5-trimethoxybenzene or ferrocene as an internal standard and *E/Z* ratios were determined using ¹H NMR of unpurified products. All other reactions were set up using standard Schlenk technique and heated with stirring in temperature controlled oil baths. **Note:** Any product yields listed in the main text that do not match those listed in the supporting information are the average of multiple isolated yields. The *E/Z* ratio of the isolated products may differ from the crude mixture due to enrichment during purification. Only the *E* isomer is reported for the ¹H and ¹³C NMR spectral data. The ¹³C NMR spectra may contain extra, unassigned peaks, which we attribute to the minor *Z* isomer. **Note:** The ¹³C NMR signal for carbons attached to boron did not appear in the collected spectra due to the quadruple splitting of ¹¹B.⁷ NMR data for some compounds may be reported in two different solvents to resolve overlapped ¹³C peaks.

2. Instrumentation and Chromatography:


400 MHz ¹H, 101 MHz ¹³C and 376 MHz ¹⁹F spectra were obtained on a 400 MHz FT-NMR spectrometer equipped with a Bruker CryoPlatform. 600 MHz ¹H, 151 MHz ¹³C, and 193 MHz ¹¹B spectra were obtained on a 600 MHz FT-NMR spectrometer equipped with a Bruker SMART probe. All samples were analyzed in the indicated deutero-solvent and were recorded at ambient temperatures. All chemical shifts are reported in ppm. ¹H NMR spectra were calibrated using the residual protio-signal in deutero-solvents as a standard. ¹³C NMR spectra were calibrated using the deutero-solvent as a standard. IR spectra were recorded on a Nicolet Magma-IR 560 FT-IR spectrometer as thin films on KBr plates. High resolution MS data was obtained on a Thermo Q-Exactive Orbitrap using electrospray ionization (ESI), or a Waters GCT Premier spectrometer using chemical ionization (CI) or liquid injection field desorption ionization (LIFDI). Column chromatography was performed with boric acid impregnated 40-63 μm silica gel (see below) with the eluent reported in parentheses. Analytical thin-layer chromatography (TLC) was performed on pre-coated glass plates and visualized by UV or by staining with iodine or KMnO₄.

3. Synthesis of Non-commercial Starting Materials and Reagents:


Alkenes:



(**S1**) A 500 mL round bottom flask equipped with a magnetic stir bar and rubber septum was charged with 4-dimethylaminopyridine (305 mg, 0.1 equiv), triethylamine (100 mL), dichloromethane (100 mL), and 5-hexenol (3.0 mL, 25 mmol, 1.0 equiv) and sealed under air. Triisopropylchlorosilane (6.7 mL, 31.3 mmol, 1.25 equiv) was then added dropwise via syringe and the reaction was stirred for 72 h. The reaction was opened to air, quenched with water (100 mL), and extracted with diethyl ether (3 x 100 mL). The combined organic layers were washed with 1M hydrochloric acid (100 mL), dried with MgSO₄, filtered through Celite, and concentrated *in vacuo*. The crude oil was purified by flash silica gel chromatography (hexanes) to afford **S1** as a colorless oil (5.32 g, 83%): ¹H NMR (400 MHz, CDCl₃) δ 5.82 (ddt, *J* = 16.9, 10.2, 6.6 Hz, 1H), 5.01 (dd, *J* = 17.1, 1.7 Hz, 1H), 4.94 (d, *J* = 10.2 Hz, 1H), 3.69 (t, *J* = 6.4 Hz, 2H), 2.08 (q, *J* = 7.1 Hz, 2H), 1.62 – 1.52 (m, 2H), 1.51 – 1.41 (m, 2H), 1.14 – 0.96 (m, 2H); ¹³C NMR (101 MHz, CDCl₃) δ 139.1, 114.5, 63.4, 33.8, 32.6, 25.3, 18.2, 12.2; FTIR (cm⁻¹): 2942, 2892, 2866, 1463, 1106, 1070, 1013, 994, 910, 882, 680, 658. HRMS (CI) m/z, calcd for [C₁₅H₃₃OSi]⁺: 257.2301; found: 257.2296.⁸




(**S2**) A 500 mL round bottom flask equipped with a magnetic stir bar and a rubber septum was flame dried under vacuum, cooled to rt and refilled with nitrogen. The flask was briefly opened to air, and quickly charged with sodium hydride 60% dispersion in mineral oil (3.5 g, 87.5 mmol, 1.5 equiv), and the flask was resealed. Anhydrous tetrahydrofuran (250 mL) was then added via syringe. The stirred suspension was cooled to 0 °C and 4-pentenol (6 mL, 58.1 mmol, 1.0 equiv) was added dropwise via syringe (*caution: vigorous gas evolution*). The cooling bath was removed, and the reaction was allowed to stir for 3 h at rt. The mixture was then recooled to 0 °C and 1-bromobutane (12.5 mL, 116.2 mmol, 2 equiv) was added dropwise via syringe. The cooling bath was removed, and the mixture was stirred overnight at rt. The reaction was opened to air and quenched with saturated ammonium chloride (50 mL), adding the first few mL dropwise. This mixture was diluted with 200 mL diethyl ether, and the organic layer was separated, dried with MgSO₄, filtered through Celite and concentrated *in vacuo*. The crude oil was purified by flash silica gel chromatography (5 : 95 ethyl acetate : hexanes) affording **S2** as a colorless oil (70%): ¹H NMR (400 MHz, CDCl₃) δ 5.82 (ddt, *J* = 16.9, 10.2, 6.6 Hz, 1H), 5.02 (dd, *J* = 17.1, 1.9 Hz, 1H), 4.95 (dd, *J* = 10.2, 2.0 Hz, 1H), 3.40 (td, *J* = 6.6, 2.8 Hz, 4H), 2.11 (q, *J* = 6.9 Hz, 2H), 1.75 – 1.62 (m, 2H), 1.59 – 1.49 (m, 2H), 1.43 – 1.31 (m, 2H), 0.91 (t, *J* = 7.4 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ 138.5, 114.8, 70.8, 70.3, 32.0, 30.5, 29.1, 19.5, 14.1; FTIR (cm⁻¹): 2954, 2924, 2854, 1457, 1377. HRMS (CI) m/z, calcd for [C₉H₁₉O]⁺: 143.1436; found: 143.1425.




(**S3**) A 250 mL 3-neck round bottom flask equipped with a magnetic stir bar and a water condenser was charged with magnesium turnings (1.4 equiv), sealed with 3 rubber septa, flame dried under vacuum, cooled to rt and refilled with nitrogen. Anhydrous diethyl ether (5 mL) was added to cover the magnesium turnings and 5-bromopentene (1.2 equiv) was added dropwise via syringe. After the solution began to self-reflux the remaining diethyl ether (95 mL) and 5-bromopentene were slowly added as needed to maintain a gentle reflux. After all of the bromide was added, the reaction was heated to 40 °C to reflux for 1 h. A separate oven dried round bottom flask equipped with a magnetic stir bar and rubber septum was cooled under vacuum, refilled with nitrogen, charged with dimethylphenylchlorosilane (2.5 mL, 15 mmol, 1.0 equiv), diluted with anhydrous diethyl ether (15 mL, 1 M) and cooled to 0 °C. The Grignard reagent, prepared above, was added dropwise via syringe to the solution of dimethylphenylchlorosilane at 0 °C. After the addition was complete, catalytic ZnCl₂ (0.5 mL, 1 M in diethyl ether, 0.03 equiv) was added via syringe and the reaction was stirred for 72 h at rt under nitrogen. The reaction was cooled to 0 °C, opened to air, diluted with diethyl ether (10 mL) then slowly quenched with water (15 mL). The mixture was extracted with diethyl ether (3 x 10 mL) and the combined organic layers were washed with brine, dried with MgSO₄, filtered through Celite, and concentrated *in vacuo*. The crude oil was purified


via flash silica gel chromatography (hexanes) to give **S3** as a clear oil (2.09 g, 68%): ¹H NMR (400 MHz, CDCl₃) δ 7.56 – 7.49 (m, 2H), 7.41 – 7.33 (m, 3H), 5.79 (ddt, J = 17.0, 10.2, 6.7 Hz, 1H), 5.06 – 4.91 (m, 2H), 2.08 (q, J = 7.1 Hz, 2H), 1.51 – 1.36 (m, 2H), 0.85 – 0.73 (m, 2H), 0.28 (s, 6H); ¹³C NMR (101 MHz, CDCl₃) δ 139.6, 139.0, 133.7, 128.9, 127.9, 114.7, 37.8, 23.5, 15.4, -2.9; FTIR (cm⁻¹): 3069, 2955, 2922, 1427, 1248, 1114, 910, 834, 811, 771, 728, 699, 469. HRMS (CI) m/z, calcd for [C₁₃H₂₁Si]⁺: 205.1413; found: 205.1429.⁹



(S4) A 100 mL round bottom flask equipped with a magnetic stir bar was charged with 4-pentenyl-1-boronic acid (1.65 g, 14.5 mmol, 1.0 equiv), pinacol (1.75 g, 14.5 mmol, 1.0 equiv), MgSO₄ (8.7 g, 5 equiv) and diluted with diethyl ether (15 mL, 1 M) under air. The reaction vessel was sealed with a rubber septum and stirred at rt overnight. The crude reaction mixture was filtered through Celite and concentrated *in vacuo*. The crude oil was purified via flash silica gel chromatography (5 : 95 ethyl acetate : hexanes) to give **S4** as a colorless oil (2.10 g, 75%): ¹H NMR (600 MHz, CDCl₃) δ 5.80 (ddt, J = 17.0, 10.2, 6.7 Hz, 1H), 4.99 (dd, 1H), 4.93 (d, 1H), 2.05 (q, J = 7.0 Hz, 2H), 1.51 (p, J = 7.7 Hz, 2H), 1.24 (s, 12H), 0.79 (t, J = 7.9 Hz, 2H); ¹³C NMR (151 MHz, CDCl₃) δ 139.1, 114.6, 83.0, 36.5, 25.0, 23.6; ¹¹B NMR (193 MHz, CDCl₃) δ 34.1; FTIR (cm⁻¹): 2979, 2931, 1407, 1378, 1318, 1146, 969, 910. HRMS (CI) m/z, calcd for [C₁₁H₂₂BO₂]⁺: 197.1713; found: 197.1701.¹⁰




(S5) A 100 mL round bottom flask equipped with a magnetic stir bar and a rubber septum was flame dried under vacuum, cooled to rt and refilled with nitrogen. Diisopropylamine (3.08 mL, 24 mmol, 1.2 equiv) and anhydrous tetrahydrofuran (10 mL) were added via syringe and the mixture was cooled to -78 °C under a positive pressure of nitrogen. n-Butyllithium (9.2 mL, 2.6 M in hexanes, 1.95 equiv) was added dropwise via syringe and the solution was stirred for 1 h at -78 °C. A solution of methyl isobutyrate (2.3 mL, 20 mmol, 1.0 equiv) in anhydrous tetrahydrofuran (10 mL, 2 M) was added dropwise via syringe and the reaction was stirred for 2 h at -78 °C. 5-Bromopentene (2.86 mL, 24 mmol, 1.2 equiv) was added via syringe then the cooling bath was removed and the reaction was stirred at rt for 24 h. The reaction was then cooled to 0 °C, opened to air and quenched with saturated aqueous ammonium chloride (20 mL). The mixture was extracted with diethyl ether (3 x 10 mL) and the combined organic layers were washed with brine, dried with MgSO₄, filtered through Celite, and concentrated *in vacuo*. The crude oil was purified via flash silica gel chromatography (5 : 95 ethyl acetate : hexanes) to give **S5** as a colorless oil (1.62 g, 48%): ¹H NMR (600 MHz, CDCl₃) δ 5.78 (ddt, J = 16.9, 10.2, 6.6 Hz, 1H), 4.99 (dd, J = 17.1 Hz, 1H), 4.94 (d, J = 10.9 Hz, 1H), 3.65 (s, 3H), 2.02 (q, J = 7.3 Hz, 2H), 1.55 – 1.47 (m, 2H), 1.39 – 1.27 (m, 2H), 1.16 (s, 6H); ¹³C NMR (151 MHz, CDCl₃) δ 178.6, 138.7, 114.7, 51.8, 42.4, 40.4, 34.2, 25.3, 24.4; FTIR (cm⁻¹): 2978, 2946, 1734, 1474, 1268, 1195, 1150, 911. HRMS (CI) m/z, calcd for [C₁₀H₁₉O₂]⁺: 171.1385; found: 171.1365.¹¹



(S6) A 500 mL round bottom flask equipped with a magnetic stir bar and a rubber septum, was flame dried under vacuum, cooled to rt, and refilled with nitrogen. Anhydrous dichloromethane (200 mL), 5-hexenol (4.8 mL, 40 mmol, 1.0 equiv), pivaloyl chloride (5.9 mL, 48 mmol, 1.2 equiv) and triethylamine (6.7 mL, 48 mmol, 1.2 equiv) were each added sequentially via syringe. The reaction was stirred at rt for 20 h. The reaction was then opened to air and quenched with 3M aqueous ammonium hydroxide (60 mL). The organic layer was separated then washed with an additional 3 M ammonium hydroxide (1 x 60 mL) then with brine (1 x 60 mL). The combined aqueous layers were extracted with dichloromethane (1 x 60 mL). The combined organic layers were dried with MgSO₄, filtered through Celite, and concentrated *in vacuo*. The crude liquid was purified via flash silica gel chromatography (5 : 95 diethyl ether : hexanes) to give **S6** as a colorless liquid (6.57g, 89%): ¹H NMR (400 MHz, CDCl₃) δ 5.79 (ddt, J = 16.9, 10.2, 6.7 Hz, 1H), 5.01 (dd, J = 17.1, 1.6 Hz, 1H), 4.96 (dd, J = 10.2, 3.1, 1.2 Hz, 1H), 4.05 (t, J = 6.6 Hz, 2H), 2.08 (q, J = 7.2 Hz, 2H), 1.74 – 1.57 (m, 2H), 1.54 – 1.37 (m, 2H), 1.19 (s, 9H); ¹³C NMR (101 MHz, CDCl₃) δ 178.8, 138.6, 114.9,

64.4, 38.9, 33.4, 28.2, 27.3, 25.3; FTIR (cm^{-1}): 2975, 2936, 1730, 1481, 1285, 1156, 911. HRMS (Cl) m/z, calcd for $[\text{C}_{11}\text{H}_{21}\text{O}_2]^+$: 185.1542; found: 185.1525.

Single Component Catalyst ($(\text{L}1)_2\text{PdCl}_2$):



$(\text{L}1)_2\text{PdCl}_2$ Bis(acetonitrile)dichloropalladium(II) (311 mg, 1.2 mmol, 1.0 equiv) and bis(3,5-di-*tert*-butylphenyl)(*tert*-butyl)phosphine (1.12 g, 2.4 mmol, 2.0 equiv) was added to a 50 mL round bottom flask equipped with a magnetic stir bar and a rubber septum. The atmosphere in the flask was replaced with nitrogen, and anhydrous dichloromethane (20 mL) was added via syringe. The reaction was stirred for 30 minutes at rt. The flask was then briefly opened to air and the solvent was removed in *vacuo* (~200 mtorr). The resulting solid

was recrystallized from anhydrous diethyl ether (15 mL) under a nitrogen atmosphere at 0 °C for 24 h to give $(\text{L}1)_2\text{PdCl}_2$ as an air stable, fine yellow solid (1.23 g, 92% yield): ^1H NMR (400 MHz, C_6D_6) δ 8.08 (t, J = 5.1 Hz, 8H), 7.60 (s, 4H), 1.76 (t, J = 7.3 Hz, 18H), 1.33 (s, 72H); ^{13}C NMR (101 MHz, C_6D_6) δ 149.6 (t, J = 4.7 Hz), 131.0 (t, J = 5.8 Hz), 130.5 (t, J = 40.1 Hz), 123.8, 36.6 (t, J = 10.6 Hz), 35.2, 31.6, 31.1 (t, J = 2.6 Hz); ^{31}P NMR (162 MHz, C_6D_6) δ 41.9; FTIR (cm^{-1}): 2962, 2903, 2868, 1477, 1419, 1362, 1248, 1135, 708; mp = 261 °C (dec.). HRMS (LIFDI) m/z, calcd for $[\text{C}_{64}\text{H}_{102}\text{Cl}_2\text{P}_2\text{Pd}]$: 1108.5885; found: 1108.5583. Anal. Calcd for $\text{C}_{64}\text{H}_{102}\text{Cl}_2\text{P}_2\text{Pd}$: C, 69.20; H, 9.26; Cl, 6.38. Found: C, 69.04; H, 9.38; Cl, 6.23.

A small portion of $(\text{L}1)_2\text{PdCl}_2$ was dissolved in dichloromethane under air and recrystallized via slow evaporation at rt to give an X-ray quality crystal (Figure S1, see below for full crystallographic details).





Figure S1. Crystal Structure of $(\text{L}1)_2\text{PdCl}_2$ (hydrogen atoms omitted for clarity).

Catecholchloroborane (catBCI):

Note: Although catecholchloroborane is commercially available, for this communication we prepared it using a modification of literature procedures.^{12,13}

Caution: This reaction generates 2 equivalents of anhydrous hydrogen chloride gas. The addition rate of boron trichloride must be controlled to prevent over-pressurization of the reaction vessel. All gas leaving the reaction should be scrubbed by base to prevent harm to people or equipment.



(catBCI) A flame dried 500 mL Schlenk flask equipped with a magnetic stir bar and a rubber septum was cooled to rt under vacuum and refilled with nitrogen. The flask was briefly opened and quickly charged with catechol (10.46 g, 95 mmol, 0.95 equiv) then resealed. The flask was then evacuated and refilled with nitrogen 3 times. Anhydrous hexanes (200 mL) was added via canula transfer. With a positive flow of nitrogen, the septum was removed and the Schlenk flask was equipped with an oven dried 100 mL pressure-equalizing addition funnel equipped with a rubber septum. The apparatus was fitted with a vent tube attached to a gas scrubber filled with aqueous NaOH (250 mL, ~10 M, 0 °C). The catechol solution was cooled to 0 °C with stirring (~1 h). Boron trichloride (100 mL, 1.0 M in hexane, 1.0 equiv) was cannula transferred into the addition funnel (stopcock closed). The stopcock was opened slightly to add boron trichloride dropwise to the solution of catechol with rapid stirring. Once the addition of boron trichloride was complete, the reaction was allowed to warm to rt and stir overnight (8-12 h) with a slow, constant flow of nitrogen entering the Schlenk flask through the side arm, and exiting the addition funnel through the vent tube leading to the gas scrubber. After 8-12 h, when the solution has become homogeneous, the slow addition funnel was quickly replaced with a glass stopper and the hexanes was removed *in vacuo* (~200 mtorr) through a solid NaOH trap to quench any residual hydrogen chloride gas. Once dried, the flask was sealed under vacuum, brought into a dry glovebox and scraped into vials. The catecholchloroborane was used without further purification, but it can be sublimed (70 °C, 20 torr) to give a white solid (13.77 g, 94%): ¹H NMR (400 MHz, CDCl₃) δ 7.18 – 7.12 (m, 2H), 7.07 – 7.00 (m, 2H); ¹³C NMR (101 MHz, CDCl₃) δ 148.2, 123.6, 112.9; ¹¹B NMR (128 MHz, CDCl₃) δ 28.9.

Comments and Preparation of Boric Acid Impregnated Silica:

Note: Many boronic esters are known to partially decompose during purification on silica gel. However, boric acid impregnated silica has been shown to decrease this decomposition resulting in increased isolated yields.¹⁴ In our hands, use of commercial, untreated silica gel resulted in yields ~5-10 % lower than with boric acid impregnated silica gel (prepared according to the procedure below). Importantly, while boric acid impregnated silica gel is not necessary for this method, the use of it for purification did result in more consistent isolated yields.

Boric Acid Impregnated Silica: A 1000 mL round bottom flask equipped with a thick stir bar was charged with silica gel (300 mL), boric acid (28 g) and ethanol (550 mL) under air. The suspension was stirred at rt for 2 h at which time the ethanol and excess boric acid were removed by filtration through a fine glass frit. The impregnated silica was washed with ethanol (3 x 200 mL) and left to dry on the frit under constant suction for 16 h then placed in a 140 °C gravity oven to dry for an additional 48 h. Once dried, the boric acid impregnated silica was stored at rt in a desiccator until used.


4. General Procedure for the α -Olefin Boryl-Heck Reaction:

Note: All reactions in this section were performed on a 3 mmol scale.


General Procedure A (α -Olefins):

In a nitrogen-filled glovebox, catecholchloroborane (1.5 equiv), lithium trifluoromethanesulfonate (1.5 equiv) and $(\text{L}1)_2\text{PdCl}_2$ (2.5 mol %), were added to a flame-dried or oven-dried 25 mL Schlenk flask equipped with a stir bar. The flask was sealed with a rubber septum, removed from the glovebox and attached to a nitrogen manifold via the sidearm. 3 mL of trifluorotoluene (1 mL per mmol) was added via syringe with stirring. N,N-dicyclohexylmethylamine (5.0 equiv) was added via syringe and then the sides of the flask were rinsed with 3 mL of additional trifluorotoluene (1 mL per mmol). The solution was stirred in an oil bath at 90 °C for 15 minutes, at which time the alkene substrate (1.0 equiv) was added in one portion via syringe. The reaction was allowed to stir at 90 °C for 24 h. After 24 h, the reaction was removed from the oil bath and opened to air. Pinacol (3 equiv) was added in one portion and the reaction was removed from heat and stirred for 1 h at rt. At that time, and with the reactor now cooled to rt, the reaction was diluted with 20 mL of diethyl ether. The reaction was stirred for 10 min, filtered through Celite and concentrated *in vacuo* to remove solvents, including trifluorotoluene. The crude oil was diluted with diethyl ether (30 mL) and washed with 1 M hydrochloric acid (3 x 20 mL) to remove excess amine. The organic layer was stirred with ammonium pyrrolidine-dithiocarbamate (palladium scavenger, 6 equiv to palladium)¹⁵ for 1 h, then dried with MgSO₄, filtered through Celite and concentrated *in vacuo*. A small aliquot of the crude reaction mixture (~20 μL) was analyzed by ¹H NMR to determine the E/Z ratio. The crude material was purified via silica column chromatography on boric acid impregnated silica gel (as prepared above) in the indicated solvent combination.

Note: Only the E isomer is reported for the ¹H and ¹³C NMR spectral data. The ¹³C NMR spectra may contain extra, unassigned peaks, which are attributed to the minor Z isomer.




(5) According to general procedure A, catecholchloroborane (694 mg, 4.5 mmol), $(\text{L}1)_2\text{PdCl}_2$ (83 mg, 0.075 mmol), lithium trifluoromethanesulfonate (702 mg, 4.5 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and 1-decene (443 mg, 3.0 mmol) were combined under N₂ and stirred at 90 °C for 24 h. The reaction was worked up according to general procedure A. Analysis of the crude reaction mixture via ¹H NMR revealed a 88:12 E/Z ratio. The product was purified via boric acid impregnated silica gel chromatography (5 : 95 dichloromethane : hexanes) to afford **5** as a colorless oil (747 mg, 94%): ¹H NMR (600 MHz, CDCl₃) δ 6.64 (dt, *J* = 17.9, 6.5 Hz, 1H), 5.42 (dd, *J* = 17.9, 1.9 Hz, 1H), 2.20 – 2.08 (m, 2H), 1.45 – 1.36 (m, 2H), 1.30 – 1.22 (m, 22H), 0.87 (t, 3H); ¹H NMR (400 MHz, C₆D₆) δ 7.00 (dt, *J* = 17.8, 6.5 Hz, 1H), 5.82 (dt, *J* = 17.8, 1.5 Hz, 1H), 2.07 (q, 2H), 1.42 – 1.13 (m, 12H), 1.10 (s, 12H), 0.92 – 0.83 (m, 3H); ¹³C NMR (151 MHz, CDCl₃) δ 155.0, 83.1, 36.0, 32.0, 29.6, 29.4, 28.4, 24.9, 22.8, 14.3; ¹³C NMR (101 MHz, C₆D₆) δ 155.1, 82.9, 36.3, 32.3, 29.9, 29.7, 29.6, 28.8, 25.0, 23.1, 14.4; ¹¹B NMR (193 MHz, CDCl₃) δ 29.9; ¹¹B NMR (128 MHz, C₆D₆) δ 30.0; FTIR (cm⁻¹): 2978, 2926, 2855, 1639, 1362, 1320, 1146. HRMS (Cl) m/z, calcd for [C₁₆H₃₂BO₂]⁺: 267.2495; found: 267.2484.




(6) According to general procedure A, catecholchloroborane (694 mg, 4.5 mmol), $(\text{L}1)_2\text{PdCl}_2$ (83 mg, 0.075 mmol), lithium trifluoromethanesulfonate (702 mg, 4.5 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and 4-phenyl-1-butene (397 mg, 3.0 mmol) were combined under N₂ and stirred at 90 °C for 24 h. The reaction was worked up according to general procedure A. Analysis of the crude reaction mixture via ¹H NMR revealed a 88:12 E/Z ratio. The product was purified via boric acid impregnated silica gel chromatography (3 : 7 dichloromethane : hexanes) to afford **6** as a colorless oil (687 mg, 89%): ¹H NMR (600 MHz, CDCl₃) δ 7.30 – 7.27 (m, 2H), 7.20 – 7.16 (m,


3H), 6.70 (dt, $J = 18.0, 6.2$ Hz, 1H), 5.50 (d, $J = 18.0$ Hz, 1H), 2.79 – 2.68 (m, 2H), 2.53 – 2.39 (m, 2H), 1.27 (s, 12H); ^1H NMR (400 MHz, C_6D_6) δ 7.18 – 6.90 (m, 6H), 5.78 (d, $J = 17.9$ Hz, 1H), 2.57 – 2.48 (m, 2H), 2.37 – 2.25 (m, 2H), 1.09 (s, 12H); ^{13}C NMR (101 MHz, CDCl_3) δ 153.6, 141.9, 128.5, 126.0, 83.2, 37.7, 34.7, 24.9; ^{11}B NMR (193 MHz, CDCl_3) δ 29.7; ^{11}B NMR (128 MHz, C_6D_6) δ 29.8; FTIR (cm^{-1}): 2977, 2929, 1637, 1361, 1320, 1144. HRMS (CI) m/z, calcd for $[\text{C}_{16}\text{H}_{24}\text{BO}_2]^+$: 259.1869; found: 259.1867.




(7) According to general procedure A, catecholchloroborane (694 mg, 4.5 mmol), $(\text{L1})_2\text{PdCl}_2$ (83 mg, 0.075 mmol), lithium trifluoromethanesulfonate (702 mg, 4.5 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and vinylcyclohexane (341 mg, 3.0 mmol) were combined under N_2 and stirred at 90 °C for 24 h. The reaction was worked up according to general procedure A. Analysis of the crude reaction mixture via ^1H NMR revealed a 96:4 E/Z ratio. The product was purified via boric acid impregnated silica gel chromatography (1 : 9 dichloromethane : hexanes) to afford 7 as a pale yellow oil (595 mg, 84%): ^1H NMR (600 MHz, CDCl_3) δ 6.58 (dd, $J = 18.2, 6.2$ Hz, 1H), 5.38 (d, $J = 18.2$, 1H), 2.07 – 1.94 (m, 1H), 1.80 – 1.67 (m, 4H), 1.67 – 1.60 (m, 2H), 1.27 (s, 12H), 1.20 – 1.00 (m, 4H); ^{13}C NMR (151 MHz, CDCl_3) δ 160.0, 83.1, 43.4, 32.1, 26.3, 26.1, 24.9; ^{11}B NMR (193 MHz, CDCl_3) δ 30.1; FTIR (cm^{-1}): 2978, 2925, 2852, 1636, 1371, 1349, 1321, 1147. HRMS (CI) m/z, calcd for $[\text{C}_{14}\text{H}_{26}\text{BO}_2]^+$: 237.2026; found: 237.2041.



(8) According to general procedure A, catecholchloroborane (694 mg, 4.5 mmol), $(\text{L1})_2\text{PdCl}_2$ (83 mg, 0.075 mmol), lithium trifluoromethanesulfonate (702 mg, 4.5 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and 1,7-octadiene (165 mg, 1.5 mmol) were combined under N_2 and stirred at 90 °C for 24 h. The reaction was worked up according to general procedure A. Analysis of the crude reaction mixture via ^1H NMR revealed a 90:10 E/Z ratio. The product was purified via boric acid impregnated silica gel chromatography (1 : 1 dichloromethane : hexanes) to afford 8 as a white solid (450 mg, 83%): ^1H NMR (600 MHz, CDCl_3) δ 6.61 (dt, $J = 17.9, 6.5$ Hz, 2H), 5.42 (d, $J = 17.9$ Hz, 2H), 2.20 – 2.07 (m, 4H), 1.43 (p, $J = 3.5$ Hz, 4H), 1.26 (s, 24H); ^{13}C NMR (151 MHz, CDCl_3) δ 154.6, 83.0, 35.8, 27.9, 24.9; ^{11}B NMR (193 MHz, CDCl_3) δ 29.9; FTIR (cm^{-1}): 2978, 2929, 1638, 1362, 1320, 1146; mp = 53 °C. HRMS (CI) m/z, calcd for $[\text{C}_{20}\text{H}_{37}\text{B}_2\text{O}_4]^+$: 363.2878; found: 363.2879.




(9) According to general procedure A, catecholchloroborane (694 mg, 4.5 mmol), $(\text{L1})_2\text{PdCl}_2$ (83 mg, 0.075 mmol), lithium trifluoromethanesulfonate (702 mg, 4.5 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and triisopropylsiloxo-5-hexene (770 mg, 3.0 mmol) were combined under N_2 and stirred at 90 °C for 24 h. The reaction was worked up according to general procedure. Analysis of the crude reaction mixture via ^1H NMR revealed a 90:10 E/Z ratio. The product was purified via boric acid impregnated silica gel chromatography (3 : 7 dichloromethane : hexanes) to afford 9 as a colorless oil (1.02 g, 90%): ^1H NMR (600 MHz, CDCl_3) δ 6.63 (dt, $J = 17.9, 6.4$ Hz, 1H), 5.43 (d, $J = 17.9$ Hz, 1H), 3.67 (t, $J = 6.3$ Hz, 2H), 2.17 (q, 2H), 1.67 – 1.43 (m, 4H), 1.26 (s, 12H), 1.05 (s, 21H); ^{13}C NMR (151 MHz, CDCl_3) δ 154.7, 83.1, 63.3, 35.8, 32.7, 24.9, 24.7, 18.2, 12.2; ^{11}B NMR (193 MHz, CDCl_3) δ 29.8; FTIR (cm^{-1}): 2941, 2866, 1638, 1363, 1320, 1147, 1108. HRMS (CI) m/z, calcd for $[\text{C}_{21}\text{H}_{44}\text{BO}_3\text{Si}]^+$: 383.3153; found: 383.3134.




(10) According to general procedure A, catecholchloroborane (694 mg, 4.5 mmol), $(\text{L1})_2\text{PdCl}_2$ (83 mg, 0.075 mmol), lithium trifluoromethanesulfonate (702 mg, 4.5 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and 5-butoxy-pent-1-ene (426 mg, 3.0 mmol) were combined under N_2 and stirred at 90 °C for 24 h. The reaction was worked up according to general procedure A. Analysis of the crude reaction


mixture via ^1H NMR revealed a 87:13 *E/Z* ratio. The product was purified via boric acid impregnated silica gel chromatography (1 : 9 dichloromethane : hexanes) to afford **10** as a colorless oil (741 mg, 92%): ^1H NMR (600 MHz, CDCl_3) δ 6.63 (dt, J = 17.9, 6.4 Hz, 1H), 5.45 (d, J = 18.0 Hz, 1H), 3.50 – 3.30 (m, 4H), 2.28 – 2.13 (m, 2H), 1.75 – 1.63 (m, 2H), 1.54 (ddd, J = 14.6, 11.2, 4.9 Hz, 2H), 1.42 – 1.31 (m, 2H), 1.26 (s, 12H), 0.91 (t, J = 7.4 Hz, 3H); ^{13}C NMR (151 MHz, CDCl_3) δ 154.0, 83.2, 70.8, 70.3, 32.5, 32.0, 28.4, 24.9, 19.5, 14.1; ^{11}B NMR (193 MHz, CDCl_3) δ 29.8; FTIR (cm^{-1}): 2977, 2934, 2862, 1639, 1364, 1321, 1146. HRMS (CI) m/z, calcd for $[\text{C}_{15}\text{H}_{30}\text{BO}_3]^+$: 269.2288; found: 269.2274.



(**11**) According to general procedure A, catecholchloroborane (694 mg, 4.5 mmol), $(\text{L1})_2\text{PdCl}_2$ (83 mg, 0.075 mmol), lithium trifluoromethanesulfonate (702 mg, 4.5 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and 11-chloro-1-undecene (584 mg, 3.0 mmol) were combined under N_2 and stirred at 90 °C for 24 h. The reaction was worked up according to general procedure A. Analysis of the crude reaction mixture via ^1H NMR revealed a 88:12 *E/Z* ratio. The product was purified via boric acid impregnated silica gel chromatography (1 : 4 dichloromethane : hexanes) to afford **11** as a colorless oil (830 mg, 88%): ^1H NMR (600 MHz, CDCl_3) δ 6.63 (dt, J = 18.0, 6.4 Hz, 1H), 5.42 (d, J = 18.0, 1H), 3.53 (t, J = 6.8 Hz, 2H), 2.19 – 2.03 (m, 2H), 1.76 (p, 2H), 1.46 – 1.36 (m, 4H), 1.28 (s, 8H), 1.26 (s, 12H); ^1H NMR (400 MHz, C_6D_6) δ 7.01 (dt, J = 17.8, 6.5 Hz, 1H), 5.83 (d, J = 17.8 Hz, 1H), 3.12 (t, J = 6.7 Hz, 2H), 2.09 (q, J = 6.6 Hz, 2H), 1.56 – 1.23 (m, 6H), 1.21 – 0.97 (m, 20H); ^{13}C NMR (151 MHz, CDCl_3) δ 154.9, 83.1, 45.3, 36.0, 32.8, 29.5, 29.3, 29.0, 28.3, 27.0, 25.0; ^{13}C NMR (101 MHz, C_6D_6) δ 154.9, 83.0, 45.0, 36.2, 32.9, 29.71, 29.65, 29.5, 29.1, 28.8, 27.1, 25.0; ^{11}B NMR (193 MHz, CDCl_3) δ 30.0; ^{11}B NMR (128 MHz, C_6D_6) δ 29.3; FTIR (cm^{-1}): 2978, 2928, 2855, 1639, 1362, 1330, 1146. HRMS (CI) m/z, calcd for $[\text{C}_{17}\text{H}_{33}\text{BO}_2\text{Cl}]^+$: 315.2262; found: 315.2263.




(**12**) According to general procedure A, catecholchloroborane (694 mg, 4.5 mmol), $(\text{L1})_2\text{PdCl}_2$ (83 mg, 0.075 mmol), lithium trifluoromethanesulfonate (702 mg, 4.5 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and (dimethyl)pent-4-enyl(phenyl)silane (613 mg, 3.0 mmol) were combined under N_2 and stirred at 90 °C for 24 h. The reaction was worked up according to general procedure A. Analysis of the crude reaction mixture via ^1H NMR revealed a 89:11 *E/Z* ratio. The product was purified via boric acid impregnated silica gel chromatography (1 : 9 dichloromethane : hexanes) to afford **12** as a yellow oil (881 mg, 89%): ^1H NMR (400 MHz, CDCl_3) δ 7.59 – 7.44 (m, 2H), 7.40 – 7.31 (m, 3H), 6.60 (dt, J = 17.9, 6.5 Hz, 1H), 5.46 – 5.37 (m, 1H), 2.24 – 2.09 (m, 2H), 1.54 – 1.36 (m, 2H), 1.26 (s, 12H), 0.82 – 0.71 (m, 2H), 0.25 (s, 6H); ^{13}C NMR (101 MHz, CDCl_3) δ 154.6, 139.5, 133.7, 128.9, 127.8, 83.1, 39.8, 24.9, 22.9, 15.6, -2.9; ^{11}B NMR (193 MHz, CDCl_3) δ 29.8; FTIR (cm^{-1}): 2978, 2924, 1638, 1427, 1398, 1389, 1362, 1320, 1249, 1214, 1146, 972, 850, 835, 812, 729, 700. HRMS (CI) m/z, calcd for $[\text{C}_{19}\text{H}_{32}\text{BO}_2\text{Si}]^+$: 331.2265; found: 331.2245.




(**13**) According to general procedure A, catecholchloroborane (694 mg, 4.5 mmol), $(\text{L1})_2\text{PdCl}_2$ (83 mg, 0.075 mmol), lithium trifluoromethanesulfonate (702 mg, 4.5 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and pent-4-enylpinicoborane (570 mg, 2.9 mmol) were combined under N_2 and stirred at 90 °C for 24 h. The reaction was worked up according to general procedure A. Analysis of the crude reaction mixture via ^1H NMR revealed a 89:11 *E/Z* ratio. The product was purified via boric acid impregnated silica gel chromatography (1 : 9 dichloromethane : hexanes) to afford **13** as a yellow oil (677 mg, 72%): ^1H NMR (400 MHz, CDCl_3) δ 6.59 (dt, J = 17.9, 6.5 Hz, 1H), 5.39 (d, J = 17.9 Hz, 1H), 2.25 – 2.01 (m, 2H), 1.68 – 1.38 (m, 2H), 1.23 (s, 12H), 1.21 (s, 12H), 0.76 (t, J = 7.9 Hz, 2H); ^1H NMR (400 MHz, CD_3CN) δ 6.52 (dt, J = 17.9, 6.6 Hz, 1H), 5.35 (dt, J = 17.9, 1.5 Hz, 1H), 2.15 – 2.09 (m, 2H), 1.47 (p, J = 7.6 Hz, 2H), 1.21 (s, 12H), 1.20 (s, 12H), 0.70 (t, J = 7.7 Hz, 2H); ^{13}C NMR (101 MHz, CDCl_3) δ 154.7, 83.1, 83.0, 38.5, 25.0, 22.9; ^{13}C NMR (101 MHz, CD_3CN) δ 155.2, 83.8, 83.7, 38.8, 25.12, 25.07, 23.6; ^{11}B NMR (193 MHz, CDCl_3) δ 34.1, 29.9; ^{11}B NMR (128 MHz,

CD_3CN) δ 35.5, 31.2; FTIR (cm^{-1}): 2978, 2931, 1631, 1371, 1362, 1319, 1267, 1239, 1214, 1146, 969, 849; HRMS (CI) m/z, calcd for $[\text{C}_{17}\text{H}_{33}\text{B}_2\text{O}_4]^+$: 323.2565; found: 323.2576.




(16) According to general procedure A, catecholchloroborane (694 mg, 4.5 mmol), $(\text{L1})_2\text{PdCl}_2$ (83 mg, 0.075 mmol), lithium trifluoromethanesulfonate (702 mg, 4.5 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and methyl 2,2-dimethylhept-6-enoate (511 mg, 3.0 mmol) were combined under N_2 and stirred at 90 °C for 24 h.

The reaction was worked up according to general procedure A. Analysis of the crude reaction mixture via ^1H NMR revealed a 87:13 *E/Z* ratio. The product was purified via boric acid impregnated silica gel chromatography (7 : 3 dichloromethane : hexanes) to afford **16** as a yellow oil (732 mg, 82%): ^1H NMR (400 MHz, CDCl_3) δ 6.60 (dt, $J = 18.0, 6.4$ Hz, 1H), 5.42 (dt, $J = 18.0, 1.6$ Hz, 1H), 3.65 (s, 3H), 2.17 – 1.99 (m, 2H), 1.55 – 1.46 (m, 2H), 1.34 (tdd, $J = 8.9, 7.1, 4.5$ Hz, 2H), 1.26 (s, 12H), 1.15 (s, 6H); ^{13}C NMR (101 MHz, CDCl_3) δ 178.6, 154.2, 83.2, 51.8, 42.4, 40.5, 36.3, 25.3, 24.9, 23.8; ^{11}B NMR (193 MHz, CDCl_3) δ 30.2; FTIR (cm^{-1}): 2978, 2944, 1733, 1639, 1362, 1320, 1146. HRMS (CI) m/z, calcd for $[\text{C}_{16}\text{H}_{30}\text{BO}_4]^+$: 297.2237; found: 297.2240.



(17) According to general procedure A, catecholchloroborane (694 mg, 4.5 mmol), $(\text{L1})_2\text{PdCl}_2$ (83 mg, 0.075 mmol), lithium trifluoromethanesulfonate (702 mg, 4.5 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and hex-5-enyl pivalate (553 mg, 3.0 mmol)

were combined under N_2 and stirred at 90 °C for 24 h. The reaction was worked up according to general procedure A. Analysis of the crude reaction mixture via ^1H NMR revealed a 88:12 *E/Z* ratio. The product was purified via boric acid impregnated silica gel chromatography (3 : 7 dichloromethane : hexanes) to afford **17** as a yellow oil (847 mg, 91%): ^1H NMR (600 MHz, CDCl_3) δ 6.61 (dt, $J = 18.0, 6.4$ Hz, 1H), 5.44 (d, $J = 18.0$ Hz, 1H), 4.04 (t, $J = 6.5$ Hz, 2H), 2.19 (tdd, $J = 7.7, 6.3, 1.5$ Hz, 2H), 1.68 – 1.59 (m, 2H), 1.53 – 1.44 (m, 2H), 1.26 (s, 12H), 1.19 (s, 9H); ^{13}C NMR (101 MHz, CDCl_3) δ 178.8, 154.0, 83.2, 64.3, 38.9, 35.4, 28.3, 27.3, 24.9, 24.7; ^{11}B NMR (193 MHz, CDCl_3) δ 29.8; FTIR (cm^{-1}): 2978, 2935, 1730, 1269, 1363, 1147. HRMS (CI) m/z, calcd for $[\text{C}_{17}\text{H}_{32}\text{BO}_4]^+$: 311.2394; found: 311.2379.



(18) According to general procedure A, catecholchloroborane (694 mg, 4.5 mmol), $(\text{L1})_2\text{PdCl}_2$ (83 mg, 0.075 mmol), lithium trifluoromethanesulfonate (702 mg, 4.5 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and allylbenzene (356 mg, 3.0 mmol) were combined under N_2 and stirred at 90 °C for 24 h.


The reaction was worked up according to general procedure A. The product was purified via boric acid impregnated silica gel chromatography (3 : 7 dichloromethane : hexanes) to afford four isomers of **18** as a yellow oil (527 mg, 72%): ^1H NMR (400 MHz, CDCl_3) *E*-allyl product: δ 7.35 – 7.14 (m, 5H), 6.38 (d, $J = 15.8$ Hz, 1H), 6.30 (m, 1H), 1.88 (d, $J = 7.0$ Hz, 2H), 1.26 (s, 12H), ^{11}B NMR (128 MHz, CDCl_3) δ 33.6; *E*-alkenyl product: δ 7.35 – 7.14 (m, 5H), 6.76 (dt, $J = 17.8, 6.3$ Hz, 1H), 5.45 (dt, $J = 17.9, 1.6$ Hz, 1H), 3.48 (dd, $J = 6.3, 1.4$ Hz, 2H), 1.25 (s, 12H); ^{11}B NMR (128 MHz, CDCl_3) δ 30.4; FTIR (cm^{-1}): 3026, 2978, 2931, 1637, 1496, 1449, 1361, 1325, 1271, 1214, 1167, 1144, 1109, 998, 967, 887, 851, 745, 695, 673. HRMS (CI) m/z, calcd for $[\text{C}_{15}\text{H}_{22}\text{BO}_2]^+$: 245.1713; found: 245.1722.

5. General Procedure for the Styrene Boryl-Heck Reaction:


Note: All reactions in this section were performed on a 3 mmol scale.

General Procedure B (Styrenes):


In a nitrogen-filled glovebox, catecholchloroborane (1.5 equiv) was added to a dry 25 mL Schlenk flask equipped with a stir bar. The flask was sealed with a rubber septum, removed from the glovebox and attached to a nitrogen manifold via the sidearm. The flask was briefly opened and lithium iodide (5 mol %) and $(\text{L}1)_2\text{PdCl}_2$ (2.5 mol %) were added and the flask was resealed, then evacuated and refilled with nitrogen 3 times. 3 mL of trifluorotoluene (1 mL per mmol) was added via syringe with stirring. N,N-dicyclohexylmethylamine (5.0 equiv) was added via syringe and then the sides of the flask were rinsed with 3 mL of additional trifluorotoluene (1 mL per mmol). The solution was stirred in an oil bath at 70 °C for 15 minutes, at which time the alkene substrate (1.0 equiv) was added in one portion via syringe. The reaction was allowed to stir at 70 °C for 24 h. After 24 h, the reaction was removed from the oil bath and opened to air. Pinacol (3 equiv) was added in one portion and the reaction was removed from heat and stirred for 1 h at rt. At that time, and with the reactor now cooled to rt, the reaction was diluted with 20 mL of diethyl ether. The reaction was stirred for 10 min, filtered through Celite and concentrated *in vacuo* to remove solvents, including trifluorotoluene. The crude oil was diluted with diethyl ether (30 mL) and washed with 1 M hydrochloric acid (3 x 20 mL) to remove excess amine. The organic layer was stirred with ammonium pyrrolidine-dithiocarbamate (palladium scavenger, 6 equiv to palladium)¹⁵ for 1 h, then dried with MgSO₄, filtered through Celite and concentrated *in vacuo*. The crude material was purified via silica column chromatography on boric acid impregnated silica gel (as prepared above) in the indicated solvent combination.



(**19**) According to general procedure B, catecholchloroborane (694 mg, 4.5 mmol), $(\text{L}1)_2\text{PdCl}_2$ (83 mg, 0.075 mmol), lithium iodide (20 mg, 0.15 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and 4-*tert*-butylstyrene (481 mg, 3.0 mmol) were combined under N₂ and stirred at 70 °C for 24 h. The reaction was worked up according to general procedure B and purified on boric acid impregnated silica gel chromatography (3 : 7 dichloromethane : hexanes) to afford **19** as a white solid (814 mg, 95%): ¹H NMR (600 MHz, CDCl₃) δ 7.43 (d, *J* = 8.3 Hz, 2H), 7.41 – 7.33 (m, 3H), 6.12 (d, *J* = 18.4 Hz, 1H), 1.31 (s, 21H); ¹³C NMR (151 MHz, CDCl₃) δ 152.3, 149.5, 134.9, 127.0, 125.7, 83.4, 34.9, 31.4, 25.0; ¹¹B NMR (193 MHz, CDCl₃) δ 30.6; FTIR (cm⁻¹): 2966, 1625, 1346, 1329, 1141; mp = 89–90 °C. HRMS (Cl) m/z, calcd for [C₁₈H₂₈BO₂]⁺: 287.2182; found: 287.2191.




(**20**) According to general procedure B, catecholchloroborane (694 mg, 4.5 mmol), $(\text{L}1)_2\text{PdCl}_2$ (83 mg, 0.075 mmol), lithium iodide (20 mg, 0.15 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and 2,4-dimethylstyrene (408 mg, 3.0 mmol) were combined under N₂ and stirred at 70 °C for 24 h. The reaction was worked up according to general procedure B and purified on boric acid impregnated silica gel chromatography (7 : 13 dichloromethane : hexanes) to afford **20** as a pale yellow oil (742 mg, 99%): ¹H NMR (600 MHz, CDCl₃) δ 7.62 (d, *J* = 18.3 Hz, 1H), 7.47 (d, *J* = 7.9 Hz, 1H), 6.99 (d, *J* = 8.0 Hz, 1H), 6.97 (s, 1H), 6.04 (d, *J* = 18.3 Hz, 1H), 2.39 (s, 3H), 2.30 (s, 3H), 1.31 (s, 12H); ¹³C NMR (151 MHz, CDCl₃) δ 147.1, 138.7, 136.4, 134.0, 131.3, 127.0, 125.9, 83.4, 25.0, 21.3, 19.9; ¹¹B NMR (193 MHz, CDCl₃) δ 29.6; FTIR (cm⁻¹): 2978, 1622, 1348, 1323, 1145. HRMS (Cl) m/z, calcd for [C₁₆H₂₄BO₂]⁺: 259.1869; found: 259.1876.




(**21**) According to general procedure B, catecholchloroborane (694 mg, 4.5 mmol), $(\text{L}1)_2\text{PdCl}_2$ (83 mg, 0.075 mmol), lithium iodide (20 mg, 0.15 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and 4-vinylanisole (403 mg, 3.0


mmol) were combined under N₂ and stirred at 70 °C for 24 h. The reaction was worked up according to general procedure B and purified on boric acid impregnated silica gel chromatography (1 : 1 dichloromethane : hexanes) to afford **21** as a yellow solid (765 mg, 98%): ¹H NMR (600 MHz, CDCl₃) δ 7.44 (d, J = 8.6 Hz, 2H), 7.35 (d, J = 18.4 Hz, 1H), 6.87 (d, J = 8.7 Hz, 2H), 6.01 (d, J = 18.4 Hz, 1H), 3.81 (s, 3H), 1.31 (s, 12H); ¹³C NMR (151 MHz, CDCl₃) δ 160.4, 149.2, 130.6, 128.6, 114.1, 83.4, 55.4, 25.0; ¹¹B NMR (193 MHz, CDCl₃) δ 30.3; FTIR (cm⁻¹): 2978, 1626, 1605, 1511, 1355, 1254, 1144; mp = 55 °C. HRMS (Cl) m/z, calcd for [C₁₅H₂₂BO₃]⁺: 261.1662; found: 261.1652.




(**22**) According to general procedure B, catecholchloroborane (694 mg, 4.5 mmol), (**L1**)₂PdCl₂ (83 mg, 0.075 mmol), lithium iodide (20 mg, 0.15 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and N,N-dimethylamino-4-vinylbenzene (442 mg, 3.0 mmol) were combined under N₂ and stirred at 70 °C for 24 h. The reaction was worked up according to general procedure B and purified on boric acid impregnated silica gel chromatography (1 : 9 dichloromethane : hexanes) to afford **22** as a pale yellow solid (766 mg, 94%): ¹H NMR (400 MHz, CDCl₃) δ 7.40 (d, J = 8.8 Hz, 2H), 7.33 (d, J = 18.4 Hz, 1H), 6.66 (d, J = 8.8 Hz, 2H), 5.92 (d, J = 18.3 Hz, 1H), 2.98 (s, 6H), 1.30 (s, 12H); ¹³C NMR (151 MHz, CDCl₃) δ 151.1, 149.9, 128.5, 126.1, 112.1, 83.2, 40.5, 25.0; ¹¹B NMR (193 MHz, CDCl₃) δ 29.8; FTIR (cm⁻¹): 2977, 1604, 1522, 1430, 1388, 1351, 1319, 1227, 1197, 1182, 1166, 1143, 807; mp = 83–87 °C. HRMS (ESI) m/z, calcd for [C₁₆H₂₅BNO₂]⁺: 274.1973; found: 274.1984.



(**23**) According to general procedure B, catecholchloroborane (694 mg, 4.5 mmol), (**L1**)₂PdCl₂ (83 mg, 0.075 mmol), lithium iodide (20 mg, 0.15 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and 3-vinylanisole (403 mg, 3.0 mmol) were combined under N₂ and stirred at 70 °C for 24 h. The reaction was worked up according to general procedure B and purified on boric acid impregnated silica gel chromatography (2 : 98 ethyl acetate : hexanes) to afford **23** as a yellow oil (657 mg, 84%): ¹H NMR (600 MHz, CDCl₃) δ 7.37 (d, J = 18.4 Hz, 1H), 7.28 – 7.22 (m, 1H), 7.08 (d, J = 7.6 Hz, 1H), 7.03 (s, 1H), 6.85 (dd, J = 8.2, 2.6 Hz, 1H), 6.16 (d, J = 18.4 Hz, 1H), 3.81 (s, 3H), 1.31 (s, 12H); ¹³C NMR (151 MHz, CDCl₃) δ 159.9, 149.5, 139.1, 129.7, 120.0, 115.0, 112.1, 83.5, 55.3, 25.0; ¹¹B NMR (193 MHz, CDCl₃) δ 30.3; FTIR (cm⁻¹): 2977, 1625, 1350, 1260, 1145. HRMS (Cl) m/z, calcd for [C₁₅H₂₂BO₃]⁺: 261.1662; found: 261.1644.




(**24**) According to general procedure B, catecholchloroborane (694 mg, 4.5 mmol), (**L1**)₂PdCl₂ (83 mg, 0.075 mmol), lithium iodide (20 mg, 0.15 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and ethyl 4-vinylbenzoate (529 mg, 3.0 mmol) were combined under N₂ and stirred at 70 °C for 24 h. The reaction was worked up according to general procedure B and purified on boric acid impregnated silica gel chromatography (1 : 1 dichloromethane : hexanes) to afford **24** as a pale yellow solid (778 mg, 86%): ¹H NMR (600 MHz, CDCl₃) δ 8.01 (d, J = 8.3 Hz, 2H), 7.53 (d, J = 8.2 Hz, 2H), 7.41 (d, J = 18.4 Hz, 1H), 6.27 (d, J = 18.4 Hz, 1H), 4.37 (q, J = 7.1 Hz, 2H), 1.39 (t, J = 7.1 Hz, 3H), 1.32 (s, 12H); ¹³C NMR (151 MHz, CDCl₃) δ 166.4, 148.3, 141.8, 130.7, 130.0, 127.0, 83.7, 61.1, 25.0, 14.5; ¹¹B NMR (193 MHz, CDCl₃) δ 30.2; FTIR (cm⁻¹): 2979, 1717, 1624, 1414, 1381, 1370, 1350, 1327, 1273, 1213, 1176, 1144, 1106, 970, 850, 764; mp = 79–80 °C. HRMS (Cl) m/z, calcd for [C₁₇H₂₄BO₄]⁺: 303.1768; found: 303.1771.




(**25**) According to general procedure B, catecholchloroborane (694 mg, 4.5 mmol), (**L1**)₂PdCl₂ (83 mg, 0.075 mmol), lithium iodide (20 mg, 0.15 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and 4-fluorostyrene (386 mg, 3.0 mmol) were combined under N₂ and stirred at 70 °C for 24 h. The reaction was worked up according to general procedure B and purified on boric acid impregnated silica gel chromatography (1 : 4 dichloromethane : hexanes) to afford **25** as a white solid (582 mg, 78%): ¹H NMR (600 MHz,


CDCl_3) δ 7.51 – 7.42 (m, 2H), 7.35 (d, J = 18.4 Hz, 1H), 7.02 (t, J = 8.6 Hz, 2H), 6.07 (d, J = 18.4 Hz, 1H), 1.31 (s, 12H); ^{13}C NMR (101 MHz, CDCl_3) δ 163.3 (d, J = 250 Hz), 148.3, 133.8 (d, J = 3 Hz), 128.8 (d, J = 8 Hz), 115.7 (d, J = 22 Hz), 83.5, 25.0; ^{11}B NMR (193 MHz, CDCl_3) δ 30.3; ^{19}F NMR (565 MHz, CDCl_3) δ -112.4; FTIR (cm^{-1}): 2978, 1508, 1350, 1326, 1144; mp = 62–63 °C. HRMS (CI) m/z, calcd for $[\text{C}_{14}\text{H}_{19}\text{BO}_2\text{F}]^+$: 249.1462; found: 249.1470.




(**26**) According to general procedure B, catecholchloroborane (694 mg, 4.5 mmol), (**L1**)₂PdCl₂ (83 mg, 0.075 mmol), lithium iodide (20 mg, 0.15 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and 4-chlorostyrene (416 mg, 3.0 mmol) were combined under N_2 and stirred at 70 °C for 24 h. The reaction was worked up according to general procedure B and purified on boric acid impregnated silica gel chromatography (3 : 7 dichloromethane : hexanes) to afford **26** as a white solid (745 mg, 94%): ^1H NMR (600 MHz, CDCl_3) δ 7.41 (d, 2H), 7.33 (d, J = 18.5 Hz, 1H), 7.30 (d, 2H), 6.13 (d, J = 18.4 Hz, 1H), 1.31 (s, 12H); ^{13}C NMR (151 MHz, CDCl_3) δ 148.2, 136.1, 134.8, 129.0, 128.4, 83.6, 25.0; ^{11}B NMR (193 MHz, CDCl_3) δ 30.2; FTIR (cm^{-1}): 2974, 1627, 1492, 1411, 1380, 1372, 1356, 1323, 1212, 1166, 1146, 1089, 993, 803, 640, 491; mp = 84–85 °C. HRMS (CI) m/z, calcd for $[\text{C}_{14}\text{H}_{19}\text{BO}_2\text{Cl}]^+$: 265.1167; found: 265.1170.



(**27**) According to general procedure B, catecholchloroborane (694 mg, 4.5 mmol), (**L1**)₂PdCl₂ (83 mg, 0.075 mmol), lithium iodide (20 mg, 0.15 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and 1-(toluene-4-sulfonyl)-2-vinyl-1*H*-pyrrole (893 mg, 3.0 mmol) were combined under N_2 and stirred at 70 °C for 24 h. The reaction was worked up according to general procedure B and purified on boric acid impregnated silica gel chromatography (3 : 2 dichloromethane : hexanes) to afford **27** as a white solid (906 mg, 71%): ^1H NMR (600 MHz, CDCl_3) δ 7.98 (d, J = 8.5 Hz, 1H), 7.86 (d, J = 8.1 Hz, 1H), 7.75 (d, J = 8.5 Hz, 2H), 7.69 (s, 1H), 7.46 (d, J = 18.7 Hz, 1H), 7.37 – 7.30 (m, 1H), 7.26 (s, 1H), 7.21 (d, J = 8.2 Hz, 2H), 6.20 (d, J = 18.7 Hz, 1H), 2.33 (s, 3H), 1.32 (s, 12H); ^{13}C NMR (101 MHz, CDCl_3) δ 145.3, 140.4, 135.8, 135.0, 130.1, 128.9, 127.0, 126.4, 125.2, 123.9, 121.6, 121.0, 113.8, 83.6, 25.0, 21.7; ^{11}B NMR (193 MHz, CDCl_3) δ 29.2; FTIR (cm^{-1}): 2977, 1627, 1372, 1348, 1175, 1142, 972; mp = 69 °C. HRMS (ESI) m/z, calcd for $[\text{C}_{23}\text{H}_{27}\text{BNO}_2\text{S}]^+$: 424.1748; found: 424.1765.




(**28**) According to general procedure B, catecholchloroborane (694 mg, 4.5 mmol), (**L1**)₂PdCl₂ (83 mg, 0.075 mmol), lithium iodide (20 mg, 0.15 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and ethenyl-1,3-benzodioxole (445 mg, 3.0 mmol) were combined under N_2 and stirred at 70 °C for 24 h. The reaction was worked up according to general procedure B and purified on boric acid impregnated silica gel chromatography (2 : 3 dichloromethane : hexanes) to afford **28** as a yellow solid (616 mg, 76%): ^1H NMR (600 MHz, CDCl_3) δ 7.30 (d, J = 18.3 Hz, 1H), 7.03 (s, 1H), 6.94 (d, J = 9.3 Hz, 1H), 6.77 (d, J = 8.0 Hz, 1H), 5.99 – 5.93 (m, 3H), 1.31 (s, 12H); ^{13}C NMR (151 MHz, CDCl_3) δ 149.2, 148.5, 148.2, 132.4, 122.8, 108.4, 106.0, 101.3, 83.4, 25.0; ^{11}B NMR (193 MHz, CDCl_3) δ 30.4; FTIR (cm^{-1}): 2978, 1506, 1446, 1330, 1250, 1144, 1040; mp = 83 °C. HRMS (CI) m/z, calcd for $[\text{C}_{15}\text{H}_{20}\text{BO}_4]^+$: 275.1455; found: 275.1456.



(**29**) According to general procedure B, catecholchloroborane (694 mg, 4.5 mmol), (**L1**)₂PdCl₂ (83 mg, 0.075 mmol), lithium iodide (20 mg, 0.15 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and α -methylstyrene (354 mg, 3.0 mmol) were combined under N_2 and stirred at 70 °C for 24 h. The reaction was worked up according to general procedure B and purified on boric acid impregnated silica gel chromatography (7 : 13 dichloromethane : hexanes) to afford **29** as a light orange oil (517 mg, 71%): ^1H NMR (600 MHz, CDCl_3) δ 7.50 (d, 2H), 7.32 (t, J = 7.4 Hz, 2H), 7.28 (d, 1H), 5.76 (s, 1H), 2.41 (s, 3H), 1.32 (s, 12H); ^{13}C NMR (151 MHz, CDCl_3) δ 157.9, 144.0, 128.3, 128.1, 126.0, 83.1, 25.1, 20.2; ^{11}B NMR


(193 MHz, CDCl₃) δ 30.0; FTIR (cm⁻¹): 2978, 1621, 1355, 1145. HRMS (Cl) m/z, calcd for [C₁₅H₂₂BO₂]⁺: 245.1713; found: 245.1710.

Preparation of Non-Pinacol Containing Boronic Esters:




(30) Using a modification of general procedure B, catecholchloroborane (694 mg, 4.5 mmol), (L1)₂PdCl₂ (83 mg, 0.075 mmol), lithium iodide (20 mg, 0.15 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and 4-*tert*-butylstyrene (481 mg, 3.0 mmol) were combined under N₂ and stirred at 70 °C for 24 h. After 24 h, the reaction was removed from the oil bath and opened to air.

2-Methyl-2,4-pentanediol (1.2 mL, 9.0 mmol) was added in one portion and the reaction was allowed to cool to rt with stirring. Once at rt, the reaction was diluted with 20 mL of diethyl ether. The reaction was stirred for 10 min, filtered through Celite and concentrated *in vacuo* to remove solvents including trifluorotoluene. The crude oil was diluted with diethyl ether (30 mL) and washed with 1 M hydrochloric acid (3 x 20 mL) to remove excess amine. The organic layer was dried with MgSO₄, filtered through Celite and concentrated *in vacuo*. The crude material was purified by boric acid impregnated silica gel chromatography (3 : 7 dichloromethane : hexanes) to afford **30** as a slight red solid (620 mg, 75%): ¹H NMR (400 MHz, CDCl₃) δ 7.42 (d, J = 8.3 Hz, 2H), 7.36 – 7.30 (m, 3H), 6.06 (d, J = 18.2 Hz, 1H), 4.27 (ddt, J = 12.0, 5.9, 3.0 Hz, 1H), 1.82 (dd, J = 13.9, 3.0 Hz, 1H), 1.56 (t, J = 14.0, 11.4 Hz, 1H), 1.35 (m, J = 2.6 Hz, 6H), 1.33 (s, 3H), 1.31 (s, 9H); ¹³C NMR (101 MHz, CDCl₃) δ 151.5, 146.5, 135.4, 126.8, 125.5, 71.0, 64.9, 46.1, 34.8, 31.4, 28.3, 23.4; ¹¹B NMR (128 MHz, C₆D₆) δ 25.6; mp = 70–73 °C; FTIR (cm⁻¹): 2970, 1623, 1411, 1391, 1331, 1307, 1267, 1218, 996, 815; HRMS (LIFDI) m/z, calcd for [C₁₈H₂₇BO₂]⁺: 286.2104; found: 286.2111.



(31) Using a modification of general procedure B, catecholchloroborane (694 mg, 4.5 mmol), (L1)₂PdCl₂ (83 mg, 0.075 mmol), lithium iodide (20 mg, 0.15 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and 4-*tert*-butylstyrene (481 mg, 3.0 mmol) were combined under N₂ and stirred at 70 °C for 24 h. After 24 h, the reaction was removed from the oil bath and opened to air.


2,2-Dimethyl-1,3-propanediol (937 mg, 9.0 mmol) was added in one portion and the reaction was allowed to cool to rt with stirring. Once at rt, the reaction was diluted with 20 mL of diethyl ether. The reaction was stirred for 10 min, filtered through Celite and concentrated *in vacuo* to remove solvents including trifluorotoluene. The crude oil was diluted with diethyl ether (30 mL) and washed with 1 M hydrochloric acid (3 x 20 mL) to remove excess amine. The organic layer was dried with MgSO₄, filtered through Celite and concentrated *in vacuo*. The crude material was purified by boric acid impregnated silica gel chromatography (1 : 9 ethyl acetate : hexanes) to afford **31** as a white solid (655 mg, 80%): ¹H NMR (600 MHz, CDCl₃) δ 7.43 (d, J = 8.3 Hz, 2H), 7.35 (d, J = 8.3 Hz, 2H), 7.31 (d, J = 18.3 Hz, 1H), 6.06 (d, J = 18.3 Hz), 3.70 (s, 4H), 1.31 (s, 9H), 1.01 (s, 6H); ¹³C NMR (151 MHz, CDCl₃) δ 151.9, 147.1, 135.2, 126.9, 125.6, 72.4, 34.8, 32.0, 31.4, 22.1; ¹¹B NMR (193 MHz, CDCl₃) δ 26.3; FTIR (cm⁻¹): 2962, 2872, 1622, 1513, 1476, 1313, 1256, 996, 815; mp = 79–81 °C. HRMS (Cl) m/z, calcd for [C₁₇H₂₆BO₂]⁺: 273.2026; found: 273.2026.



(32) Using a modification of general procedure B, catecholchloroborane (694 mg, 4.5 mmol), (L1)₂PdCl₂ (83 mg, 0.075 mmol), lithium iodide (20 mg, 0.15 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and 4-*tert*-butylstyrene (481 mg, 3.0 mmol) were combined under N₂ and stirred at 70 °C for 24 h. After 24 h, the reaction was removed from the oil bath and opened to air.


(1*R*,2*R*)-1,2-diphenyl-1,2-ethanediol (1.93 g, 9.0 mmol) was added in one portion and the

reaction was allowed to cool to rt with stirring. Once at rt, the reaction was diluted with 20 mL of diethyl ether. The reaction was stirred for 10 min, filtered through Celite and concentrated *in vacuo* to remove solvents including trifluorotoluene. The crude oil was diluted with diethyl ether (30 mL) and washed with 1 M hydrochloric acid (3 x 20 mL) to remove excess amine. The organic layer was stirred with ammonium pyrrolidine-dithiocarbamate (palladium scavenger, 6 equiv to palladium)¹⁵ for 1 h, then dried with MgSO₄, filtered through Celite and concentrated *in vacuo*. The crude material was purified by boric acid impregnated silica gel chromatography (1 : 4 ethyl acetate : hexanes) to afford **32** as a thick yellow oil (929 mg, 81%): ¹H NMR (600 MHz, CDCl₃) δ 7.61 (d, *J* = 18.4 Hz, 1H), 7.50 (d, *J* = 8.3 Hz, 2H), 7.43 – 7.39 (m, 6H), 7.38 – 7.33 (m, 6H), 6.32 (d, *J* = 18.4 Hz, 1H), 5.25 (s, 2H), 1.34 (s, 9H); ¹³C NMR (101 MHz, CDCl₃) δ 152.7, 151.1, 140.4, 134.7, 129.0, 128.5, 127.2, 126.0, 125.8, 86.7, 34.9, 31.4; ¹¹B NMR (193 MHz, CDCl₃) δ 30.8; FTIR (cm⁻¹): 2963, 1622, 1344, 1319, 1175, 999, 815, 761, 698. HRMS (Cl) m/z, calcd for [C₂₆H₂₈BO₂]⁺: 383.2182; found: 383.2178.



(33) Using a modification of general procedure B, catecholchloroborane (694 mg, 4.5 mmol), (L1)₂PdCl₂ (83 mg, 0.075 mmol), lithium iodide (20 mg, 0.15 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and 4-*tert*-butylstyrene (481 mg, 3.0 mmol) were combined under N₂ and stirred at 70 °C for 24 h. After 24 h, the reaction was removed from the oil bath and opened to air. 1,8-Diaminonaphthalene (1.42 g, 9.0 mmol) was added in

one portion and the reaction was allowed to cool to rt with stirring. Once at rt, the reaction was diluted with 20 mL of diethyl ether. The reaction was stirred for 10 min, filtered through Celite and concentrated *in vacuo* to remove solvents including trifluorotoluene. The crude material was purified by boric acid impregnated silica gel chromatography (3 : 7 dichloromethane : hexanes) to afford **33** as a yellow solid (664 mg, 85%): ¹H NMR (400 MHz, CDCl₃) δ 7.46 (d, *J* = 8.4 Hz, 2H), 7.41 (d, *J* = 8.4 Hz, 2H), 7.15 (d, *J* = 10.5 Hz, 1H), 7.11 (d, *J* = 7.5 Hz, 2H), 7.02 (d, *J* = 8.1 Hz, 2H), 6.37 (d, *J* = 7.3 Hz, 2H), 6.29 (d, *J* = 18.6 Hz, 1H), 5.86 (s, 2H), 1.34 (s, 9H); ¹³C NMR (101 MHz, CDCl₃) δ 152.1, 143.7, 141.3, 136.5, 134.9, 127.7, 126.7, 125.8, 120.0, 117.7, 105.9, 34.9, 31.4; ¹¹B NMR (193 MHz, CDCl₃) δ 28.9; FTIR (cm⁻¹): 2962, 1600, 1513, 1413, 818, 762; mp = 135–136 °C. HRMS (ESI) m/z, calcd for [C₂₂H₂₄BN₂]⁺: 327.2027; found: 327.2041.



(34) Using a modification of general procedure B, catecholchloroborane (694 mg, 4.5 mmol), (L1)₂PdCl₂ (83 mg, 0.075 mmol), lithium iodide (20 mg, 0.15 mmol), N,N-dicyclohexylmethylamine (3.2 mL, 5.0 mmol), trifluorotoluene (6.0 mL, 0.5 M), and 4-*tert*-butylstyrene (481 mg, 3.0 mmol) were combined under N₂ and stirred at 70 °C for 24 h. After 24 h, the reaction was removed from the oil bath and allowed to cool to rt with stirring. Once at rt, the reaction was diluted with 20 mL of diethyl ether. The reaction was stirred for 10 min, filtered through Celite and concentrated *in vacuo* to remove solvents including trifluorotoluene. The crude oil was diluted with diethyl ether (30 mL) and washed with 1 M hydrochloric acid (3 x 20 mL) to remove excess amine. The organic layer was dried with MgSO₄, filtered through Celite and concentrated *in vacuo*. The crude solid was diluted with diethyl ether (10 mL) and cooled 0 °C in an ice bath. Once at 0 °C, potassium bifluoride (1.41 g, 18.0 mmol) was added with stirring. Deionized water (3 mL) was then added dropwise over 30 minutes, at which time the ice bath was removed and the reaction was allowed to warm to rt with stirring. Once at rt, the solvent was removed *in vacuo*. The crude solid was dissolved in acetone (20 mL), filtered through Celite and concentrated *in vacuo*. The crude material was purified by dissolving in hot acetone (10 mL) and precipitating with diethyl ether (20 mL) to afford **34** as a white solid (564 mg, 70%): ¹H NMR (600 MHz, Acetone-d₆) δ 7.27 (s, 4H), 6.62 (d, *J* = 18.2 Hz, 1H), 6.29 (d, *J* = 18.0 Hz, 1H), 1.28 (s, 9H); ¹³C NMR (101 MHz, Acetone-d₆) δ 149.1, 139.2, 134.3 (q), 126.2, 125.7, 34.8, 31.7; ¹¹B NMR (193 MHz, Acetone-d₆) δ 3.0; ¹⁹F NMR (376 MHz, Acetone-d₆) δ -141.6; FTIR (cm⁻¹): 3647, 2966, 2869, 1705, 1627, 1563, 1464, 1412, 1393, 1364, 1296, 1098, 966, 855, 754, 560, 514; mp = 263–267 °C (dec.). HRMS (LIFDI) m/z, calcd for [C₁₂H₁₅BF₃K]: 266.0856; found: 266.0872.



6. Additional Optimization Data:

Note: All reactions in this section were performed on 0.25 mmol in a nitrogen-filled glovebox.

Examination of Palladium Sources:



During initial optimization it was found that many palladium precatalysts are competent in this reaction (Table S1). Using otherwise optimized reaction conditions, a more extensive survey of palladium precatalysts was conducted (Table S2). The results from those investigations are shown below.

Table S1. Examination of Alternate Palladium Sources with Cy₃P.

 2	10 mol % [Pd], 20 mol % Cy ₃ P 1.5 equiv catBCl 5 equiv Cy ₂ NMe PhCF ₃ , 80 °C, 48 h	Alkene Isomers 3	 4
Entry	[Pd] Source	% 2 ^a	% 3 ^a
1	(COD)Pd(CH ₂ SiMe ₃) ₂	0	38
2	Pd ₂ dba ₃	0	44
3	Pd(dba) ₂	0	37
4	[(allyl)PdCl] ₂	0	38

^aYields obtained by ¹H NMR.

Table S2. Examination of Alternate Palladium Sources with L1.

 2	2.5 mol % [Pd], 5 mol % L1 1.5 equiv catBCl, 1.5 equiv LiOTf 5 equiv Cy ₂ NMe PhCF ₃ , 90 °C, 24 h	Alkene Isomers 3	 4
Entry	[Pd] Source	% 2 ^a	% 3 ^a
1	Pd ₂ dba ₃	0	0
2	Pd ₂ dba ₃ ·CHCl ₃	0	0
3	Pd(dba) ₂	0	0
4	PdCl ₂	0	12
5	(MeCN) ₂ PdCl ₂	0	13
6	(cod)PdCl ₂	0	0
7	[(cinnamyl)PdCl] ₂	0	0
8	PdI ₂	0	36
9	Pd(OAc) ₂	0	78
10	(MeCN) ₄ Pd(BF ₄) ₂	0	1
11	(L1) ₂ Pd ^b	0	0
12	none	>99	0

^aYields obtained by ¹H NMR with ferrocene as an internal standard. ^bNo external ligand added.

Examination of Ligands with Pd₂dba₃:

During reaction optimization, additional ligands were examined than were discussed in the main text. The full table of results is shown below (Table S3).

Table S3. Full Table of Ligands.

Entry	Ligand (mol %)	%2 ^a	%3 ^a	%4 ^a
1	Ph ₃ P (20 mol %)	65	10	25
2	Cy ₃ P (20 mol %)	53	15	32
3	^t Bu ₃ P (20 mol %)	0	>99	0
4	dppe (10 mol %)	>99	0	0
5	dppp (10 mol %)	96	0	4
6	^t BuXantPhos (10 mol %)	93	7	0
7	SPhos (10 mol %)	0	56	44
8	^t BuXPhos (10 mol %)	0	68	32
9	^t BuPPh ₂ (20 mol %)	19	19	62
10	L1 (20 mol %)	2	16	82

^aYields obtained by ¹H NMR with ferrocene as an internal standard.

7. Computational and Experimental Mechanistic Insight:

Computational Methods:

All DFT calculations were performed with the Gaussian 09 software package.¹⁶ Optimizations of the geometries of the minima were conducted with the B3LYP¹⁷⁻¹⁹ method. The 6-31++G(d,p) basis set was applied to all atoms. Vibrational frequencies were computed at the same level of theory for each calculation to verify that the structure is an energy minimum and to evaluate the thermochemical properties. Solvent effects were computed based upon the gas-phase optimized structures using the same level of theory with the default polarized continuum solvation model (PCM) in Gaussian.²⁰ 1,2-Dichloroethane ($\epsilon = 10.13$) was used instead of trifluorotoluene ($\epsilon = 9.18$) because the Gaussian 09 program does not contain default solvent parameters for trifluorotoluene.²¹ In this communication, all the energies are discussed in terms of enthalpy.

Coordination Studies:

Density functional theory calculations (B3LYP/6-31++G(d,p)) modeling this coordination also suggest that Cy₂NMe coordinates more reversibly with catBCI (Figure S2). Examination of the ΔH of the coordination complexes compared to the free amine and catBCI show -7.2 kcal/mol with triethylamine but only -3.2 kcal/mol with Cy₂NMe. The stability of catBCI·Et₃N sequesters the catBCI, preventing it from entering the catalytic cycle.




Figure S2. DFT calculations of coordination complexes.

Computational Investigation of E/Z Selectivity:

We sought to compare some of the experimental E/Z ratios to calculated DFT energies for several substrates. All of the following calculations (B3LYP/6-31++G(d,p)) were performed at 90 °C. Ground state calculations for each of the four possible butene product isomers were performed to determine the relative ΔH. The products of 1-butene were chosen because they should be representative of all larger α-olefines. The computed and experimentally measured E-alkenyl, Z-alkenyl, E-allyl and Z-allyl products correlate very well supporting the notion that the observed ratio is thermodynamic (Table S4). The products from allylbenzene and styrene were also examined and show similar correlations (Tables S5 and S6).

Table S4. Calculated and Experimental Values for E/Z ratios of Butene Products



entry	structure	relative energy ΔH B3LYP	theoretical ratio	experimental ratio for compound 5
1	 Bcat	0.0 kcal/mol	92	89
2	 Bcat	1.9 kcal/mol	6	11
3	 Bcat	2.8 kcal/mol	2	0
4	 Bcat	4.3 kcal/mol	0	0

Table S5. Calculated and Experimental Values for E/Z ratios of Styrene Products




entry	structure	relative energy ΔH B3LYP	theoretical ratio	experimental ratio for compound 19
1	 Bcat	0.0 kcal/mol	100	100
2	 Bcat	4.9 kcal/mol	0	0

Table S6. Calculated and Experimental Values for E/Z ratios of Allyl Benzene Products

entry	structure	relative energy ΔH B3LYP	theoretical ratio	experimental ratio for compound 18
1	 Bcat	0.0 kcal/mol	56	67
2	 Bcat	2.9 kcal/mol	1	5
3	 Bcat	0.2 kcal/mol	40	25
4	 Bcat	2.1 kcal/mol	3	3

Isomerization Study:

The reference NMR spectra used to determine the isomerization of each alkene (Figure S3, A-C) and the crude NMR spectrum (**D**) are shown below. (*E*)-1-hexenyl-catecholborane was prepared according to a literature procedure.²² The control experiment, without catalyst, shows no isomerization of (*E*)-1-hexenyl-catecholborane suggesting that the isomerization is palladium catalyzed.




Figure S3. (A) ¹H NMR spectrum of (*E*)-1-hexenyl-pinacolborane, (B) ¹H NMR spectrum of *E* and *Z* isomers of 4-phenylbutene products, (C) ¹H NMR spectrum of *E* and *Z* isomers of 1-hexene products, (D) ¹H NMR spectrum of crude reaction mixture (Equation 2).

Isomerization Study Procedure:

In a nitrogen-filled glovebox, to a 1 dram vial with a stir bar, was added (**L1**)₂PdCl₂ (7.0 mg, 0.0063 mmol, 2.5 mol %), catecholchloroborane (58 mg, 0.375 mmol, 1.5 equiv), and lithium trifluoromethanesulfonate (58 mg, 0.375 mmol, 1.5 equiv) followed by 500 μ L of trifluorotoluene. N,N-dicyclohexylmethylamine (267 μ L, 1.25 mmol, 5.0 equiv) was added and the reaction was capped and stirred at 90 °C. After 10 min, 4-phenylbutene (38 mg, 0.25 mmol, 1.0 equiv) and (*E*)-1-hexenyl-catecholborane (50 mg, 0.25 mmol, 1.0 equiv) were added via micropipette and the reaction was sealed and stirred at 90 °C for 24 h. The reaction was quenched with pinacol (176 mg, 1.5 mmol, 6.0 equiv) and stirred for 1 h at rt. The crude mixture was diluted with 1 mL of diethyl ether, filtered through Celite and concentrated *in vacuo*. The *E/Z* ratio was determined via ¹H NMR of the crude reaction mixture in CDCl₃ (Figure S3, D).

8. Crystallographic Details:


X-ray Structural Analysis for $(L1)_2PdCl_2$. The crystal was mounted using viscous oil onto a plastic mesh and cooled to the data collection temperature (200 K). Data were collected on a Bruker-AXS APEX Duo CCD diffractometer. Unit cell parameters were obtained from 36 data frames, $0.3^\circ \omega$, from three different sections of the Ewald sphere. No symmetry higher than triclinic was observed and the centrosymmetric space group option yielded chemically reasonable and computationally stable results of refinement. The data-set was treated with absorption corrections based on redundant multiscan data. The structure was solved using direct methods and refined with full-matrix, least-squares procedures on F^2 .²³ The molecule is located at the inversion center.

All non-hydrogen atoms were refined with anisotropic displacement parameters. All hydrogen atoms were treated as idealized contributions. Scattering factors are contained in the SHELXTL 6.12 program library.²³ The CIF has been deposited under CCDC 1463772.


9. References:

- (1) Pangborn, A. B.; Giardello, M. A.; Grubbs, R. H.; Rosen, R. K.; Timmers, F. J. *Organometallics* **1996**, *15*, 1518.
- (2) McAtee, J. R.; Martin, S. E. S.; Ahneman, D. T.; Johnson, K. A.; Watson, D. A. *Angew. Chem. Int. Ed.* **2012**, *51*, 3663.
- (3) McAtee, J. R.; Yap, G. P. A.; Watson, D. A. *J. Am. Chem. Soc.* **2014**, *136*, 10166.
- (4) Martin, S. E. S.; Watson, D. A. *J. Am. Chem. Soc.* **2013**, *135*, 13330.
- (5) Aslam, S. N.; Stevenson, P. C.; Phythian, S. J.; Veitch, N. C.; Hall, D. R. *Tetrahedron* **2006**, *62*, 4214.
- (6) Waser, J.; Gaspar, B.; Nambu, H.; Carreira, E. M. *J. Am. Chem. Soc.* **2006**, *128*, 11693.
- (7) Wrackmeyer, B. *Prog. Nucl. Magn. Reson. Spectrosc.* **1979**, *12*, 227.
- (8) Doi, T.; Fukuyama, T.; Minamino, S.; Husson, G.; Ryu, I. *Chem. Commun.* **2006**, 1875.
- (9) Peñafiel, I.; Pastor, I. M.; Yus, M. *Tetrahedron* **2010**, *66*, 2928.
- (10) Nakamura, M.; Hara, K.; Hatakeyama, T.; Nakamura, E. *Org. Lett.* **2001**, *3*, 3137.
- (11) Itoh, T.; Matsueda, T.; Shimizu, Y.; Kanai, M. *Chem. Eur. J.* **2015**, *21*, 15955.
- (12) He, X.; Hartwig, J. F. *Organometallics* **1996**, *15*, 400.
- (13) Bettinger, H. F.; Filthaus, M.; Bornemann, H.; Oppel, I. M. *Angew. Chem. Int. Ed.* **2008**, *47*, 4744.
- (14) Hitosugi, S.; Tanimoto, D.; Nakanishi, W.; Isobe, H. *Chem. Lett.* **2012**, *41*, 972.
- (15) Gallagher, W. P.; Vo, A. *Org. Process Res. Dev.* **2015**, *19*, 1369.
- (16) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A.; Nakatsuji, H.; Caricato, M.; Li, X.; Hratchian, H. P.; Izmaylov, A. F.; Bloino, J.; Zheng, G.; Sonnenberg, J. L.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Montgomery Jr., J. A.; Peralta, J. E.; Ogliaro, F.; Bearpark, M. J.; Heyd, J.; Brothers, E. N.; Kudin, K. N.; Staroverov, V. N.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A. P.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Rega, N.; Millam, N. J.; Klene, M.; Knox, J. E.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Zakrzewski, V. G.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Dapprich, S.; Daniels, A. D.; Farkas, Ö.; Foresman, J. B.; Ortiz, J. V.; Cioslowski, J.; Fox, D. J.; Gaussian, Inc.: Wallingford, CT, USA, 2009.
- (17) Becke, A. D. *Phys. Rev. A* **1988**, *38*, 3098.
- (18) Lee, C.; Yang, W.; Parr, R. G. *Phys. Rev. B* **1988**, *37*, 785.
- (19) Becke, A. D. *J. Chem. Phys.* **1993**, *98*, 5648.
- (20) Cossi, M.; Rega, N.; Scalmani, G.; Barone, V. *J. Comput. Chem.* **2003**, *24*, 669.
- (21) Belding, L.; Chemler, S. R.; Dudding, T. *J. Org. Chem.* **2013**, *78*, 10288.
- (22) Fairlamb, I. J. S.; Morrison, L. R.; Dickinson, J. M.; Lu, F.-J.; Schmidt, J. P. *Bioorg. Med. Chem.* **2004**, *12*, 4285.
- (23) Sheldrick, G. *Acta Cryst. A* **2008**, *64*, 112.

10. Spectral Data:



Parameter	Value
1 Title	J301178.2.fid
2 Solvent	CDCl ₃
3 Temperature	298.2
4 Number of Scans	32
5 Receiver Gain	512.0
6 Relaxation Delay	3.0000
7 Pulse Width	9.2500
8 Spectrometer Frequency	100.62
9 Nucleus	¹³ C





—63.40


—114.45


—139.12


✓ 33.76
 ✓ 32.63
 ✓ 25.34
 ✓ 18.18
 ✓ 12.15

















Parameter	Value
1 Title	WBR03131F1.3.fid
2 Solvent	CDCl ₃
3 Temperature	298.0
4 Number of Scans	256
5 Receiver Gain	2050.0
6 Relaxation Delay	5.0000
7 Pulse Width	10.6300
8 Spectrometer Frequency	150.97
9 Nucleus	¹³ C





—34.10


Parameter	Value
1 Title	WBR03131F1.2.fid
2 Solvent	CDCl ₃
3 Temperature	298.0
4 Number of Scans	128
5 Receiver Gain	322.0
6 Relaxation Delay	1.0000
7 Pulse Width	11.0000
8 Spectrometer Frequency	192.61
9 Nucleus	¹¹ B


PinB
S4














Parameter	Value
1 Title	SBK01267A.11.fid
2 Solvent	C6D6
3 Temperature	298.0
4 Number of Scans	16
5 Receiver Gain	256.0
6 Relaxation Delay	1.0000
7 Pulse Width	10.7700
8 Spectrometer Frequency	600.32
9 Nucleus	¹ H





1.46
1.45
1.44
1.43
1.42
1.41
1.40
1.39
1.38
1.37
1.36
1.35
1.34
1.33
1.32
1.31
1.30
1.29
1.28
1.27
1.26
1.25
1.24
1.23
1.22
1.21
1.20
1.19
1.18
1.17
1.16
1.15
1.14
1.13
1.12
1.11
1.10
1.09
1.08
1.07
1.06
1.05
1.04
1.03
1.02
1.01
1.00
0.99
0.98
0.97
0.96
0.95
0.94
0.93
0.92
0.91
0.90
0.89
0.88
0.87
0.86
0.85
0.84
0.83
0.82
0.81
0.80
0.79
0.78
0.77
0.76
0.75
0.74
0.73
0.72
0.71
0.70
0.69
0.68
0.67
0.66
0.65
0.64
0.63
0.62
0.61
0.60
0.59
0.58
0.57
0.56
0.55
0.54
0.53
0.52
0.51
0.50
0.49
0.48
0.47
0.46
0.45
0.44
0.43
0.42
0.41
0.40
0.39
0.38
0.37
0.36
0.35
0.34
0.33
0.32
0.31
0.30
0.29
0.28
0.27
0.26
0.25
0.24
0.23
0.22
0.21
0.20
0.19
0.18
0.17
0.16
0.15
0.14
0.13
0.12
0.11
0.10
0.09
0.08
0.07
0.06
0.05
0.04
0.03
0.02
0.01
0.00







-41.86


Parameter	Value
1 Title	SBK01280B.1.fid
2 Solvent	C6D6
3 Temperature	298.2
4 Number of Scans	16
5 Receiver Gain	5160.6
6 Relaxation Delay	2.0000
7 Pulse Width	9.2200
8 Spectrometer Frequency	161.97
9 Nucleus	³¹ P








Parameter	Value
1 Title	WBR03089F1.2.fid
2 Solvent	CDCl ₃
3 Temperature	300.0
4 Number of Scans	16
5 Receiver Gain	203.0
6 Relaxation Delay	1.0000
7 Pulse Width	8.3000
8 Spectrometer Frequency	128.38
9 Nucleus	11B




—28.91





Parameter	Value
1 Title	WBR03128crd.1.fid
2 Solvent	CDCl ₃
3 Temperature	298.0
4 Number of Scans	16
5 Receiver Gain	71.8
6 Relaxation Delay	10.0000
7 Pulse Width	10.7700
8 Spectrometer Frequency	600.32
9 Nucleus	¹ H


Me  Bpin


5-crude














Parameter	Value
1 Title	WBR03128F1.2.fid
2 Solvent	CDCl ₃
3 Temperature	298.0
4 Number of Scans	128
5 Receiver Gain	362.0
6 Relaxation Delay	1.0000
7 Pulse Width	11.0000
8 Spectrometer Frequency	192.61
9 Nucleus	11B


—29.92





Parameter	Value
1 Title	SBK01282cr.1.fid
2 Solvent	CDCl ₃
3 Temperature	297.4
4 Number of Scans	16
5 Receiver Gain	228.0
6 Relaxation Delay	1.0000
7 Pulse Width	10.7700
8 Spectrometer Frequency	600.32
9 Nucleus	¹ H








1.03
1.04
1.05
1.09
2.28
2.28
2.29
2.32
2.32
2.33
2.51
2.53
2.54


5.76
5.80
6.91
6.93
6.94
6.96
6.97
7.02
7.04
7.08
7.10
7.12

Parameter	Value
1 Title	WBR03189D.1.fid
2 Solvent	C6D6
3 Temperature	301.1
4 Number of Scans	16
5 Receiver Gain	12.7
6 Relaxation Delay	1.0000
7 Pulse Width	15.0000
8 Spectrometer Frequency	400.13
9 Nucleus	1H












Parameter	Value
1 Title	WBR03189D-11B.1.fid
2 Solvent	C6D6
3 Temperature	303.0
4 Number of Scans	32
5 Receiver Gain	203.0
6 Relaxation Delay	1.0000
7 Pulse Width	8.3000
8 Spectrometer Frequency	128.38
9 Nucleus	11B





—29.79




Parameter	Value
1 Title	WBR03122crd.1.fid
2 Solvent	CDCl ₃
3 Temperature	298.0
4 Number of Scans	16
5 Receiver Gain	64.0
6 Relaxation Delay	10.0000
7 Pulse Width	10.7700
8 Spectrometer Frequency	600.32
9 Nucleus	¹ H









Parameter	Value
1 Title	WBR03126B.2.fid
2 Solvent	CDCl ₃
3 Temperature	298.0
4 Number of Scans	128
5 Receiver Gain	362.0
6 Relaxation Delay	1.0000
7 Pulse Width	11.0000
8 Spectrometer Frequency	192.61
9 Nucleus	11B





—30.05




Parameter	Value
1 Title	WBR03121crd.1.fid
2 Solvent	CDCl ₃
3 Temperature	298.0
4 Number of Scans	16
5 Receiver Gain	57.0
6 Relaxation Delay	10.0000
7 Pulse Width	10.7700
8 Spectrometer Frequency	600.32
9 Nucleus	¹ H












Parameter	Value
1 Title	SBK01278cr.10.fid
2 Solvent	CDCl ₃
3 Temperature	297.1
4 Number of Scans	16
5 Receiver Gain	144.0
6 Relaxation Delay	1.0000
7 Pulse Width	10.7700
8 Spectrometer Frequency	600.32
9 Nucleus	¹ H












-29.80


Parameter	Value
1 Title	SBK01278.11.fid
2 Solvent	CDCl ₃
3 Temperature	297.1
4 Number of Scans	128
5 Receiver Gain	322.0
6 Relaxation Delay	1.0000
7 Pulse Width	11.0000
8 Spectrometer Frequency	192.61
9 Nucleus	11B




Parameter	Value
1 Title	WBR03130crd.1.fid
2 Solvent	CDCl ₃
3 Temperature	298.2
4 Number of Scans	16
5 Receiver Gain	5.7
6 Relaxation Delay	1.0000
7 Pulse Width	15.0000
8 Spectrometer Frequency	400.13
9 Nucleus	1H












Parameter	Value
1 Title	WBR03120crd.1.fid
2 Solvent	CDCl ₃
3 Temperature	298.0
4 Number of Scans	16
5 Receiver Gain	161.0
6 Relaxation Delay	1.0000
7 Pulse Width	10.7700
8 Spectrometer Frequency	600.32
9 Nucleus	¹ H





11-crude















Parameter	Value
1 Title	WBR03120-C6D6.1.fid
2 Solvent	C6D6
3 Temperature	300.0
4 Number of Scans	128
5 Receiver Gain	203.0
6 Relaxation Delay	1.0000
7 Pulse Width	8.3000
8 Spectrometer Frequency	128.39
9 Nucleus	11B



11


-29.25





Parameter	Value
1 Title	WBR03157crd.1.fid
2 Solvent	CDCl ₃
3 Temperature	297.1
4 Number of Scans	16
5 Receiver Gain	57.0
6 Relaxation Delay	10.0000
7 Pulse Width	10.7700
8 Spectrometer Frequency	600.32
9 Nucleus	¹ H





12-crude




Parameter	Value
1 Title	WBR03157again.1.fid
2 Solvent	CDCl ₃
3 Temperature	298.1
4 Number of Scans	16
5 Receiver Gain	8.0
6 Relaxation Delay	1.0000
7 Pulse Width	15.0000
8 Spectrometer Frequency	400.13
9 Nucleus	¹ H









Parameter	Value
1 Title	SBK01281cr1.fid
2 Solvent	CDCl ₃
3 Temperature	297.6
4 Number of Scans	16
5 Receiver Gain	203.0
6 Relaxation Delay	1.0000
7 Pulse Width	10.7700
8 Spectrometer Frequency	600.32
9 Nucleus	¹ H







—154.73


✓22.93
✓24.96


—38.50

✓83.08
✓83.04

Parameter	Value
1 Title	SBK01281.2.fid
2 Solvent	CDCl ₃
3 Temperature	298.1
4 Number of Scans	32
5 Receiver Gain	512.0
6 Relaxation Delay	3.0000
7 Pulse Width	9.2500
8 Spectrometer Frequency	100.62
9 Nucleus	¹³ C


PinB  Bpin
13





Parameter	Value
1 Title	SBK01281.1.fid
2 Solvent	CDCl ₃
3 Temperature	297.4
4 Number of Scans	128
5 Receiver Gain	322.0
6 Relaxation Delay	1.00000
7 Pulse Width	11.00000
8 Spectrometer Frequency	192.61
9 Nucleus	11B



-29.86
-34.10





-31.20


-35.46


Parameter	Value
1 Title	SBK01281-acetonitrile.1.fid
2 Solvent	CD3CN
3 Temperature	299.4
4 Number of Scans	128
5 Receiver Gain	203.0
6 Relaxation Delay	1.0000
7 Pulse Width	8.3000
8 Spectrometer Frequency	128.39
9 Nucleus	11B




Parameter	Value
1 Title	WBR03133A.1.fid
2 Solvent	CDCl ₃
3 Temperature	298.0
4 Number of Scans	16
5 Receiver Gain	144.0
6 Relaxation Delay	1.0000
7 Pulse Width	10.7700
8 Spectrometer Frequency	600.32
9 Nucleus	¹ H












Parameter	Value
1 Title	WBR03125crd.1.fid
2 Solvent	CDCl ₃
3 Temperature	298.1
4 Number of Scans	16
5 Receiver Gain	10.1
6 Relaxation Delay	1.0000
7 Pulse Width	15.0000
8 Spectrometer Frequency	400.13
9 Nucleus	1H










Parameter	Value
1 Title	WBR03126G.2.fid
2 Solvent	CDCl ₃
3 Temperature	298.0
4 Number of Scans	128
5 Receiver Gain	362.0
6 Relaxation Delay	1.0000
7 Pulse Width	11.0000
8 Spectrometer Frequency	192.61
9 Nucleus	11B




—29.83










Parameter	Value
1 Title	SBK02001.2.fid
2 Solvent	DCI β
3 Temperature	300.0
4 Number of Scans	128
5 Receiver Gain	203.0
6 Relaxation Delay	1.0000
7 Pulse Width	8.3000
8 Spectrometer Frequency	128.39
9 Nucleus	11B



—33.56
—30.43



Parameter	Value
1 Title	SBK01253P1.fid
2 Solvent	CDCl ₃
3 Temperature	298.0
4 Number of Scans	16
5 Receiver Gain	161.0
6 Relaxation Delay	1.0000
7 Pulse Width	10.7700
8 Spectrometer Frequency	600.32
9 Nucleus	¹ H





✓24.98
✓31.40
✓34.85

-83.42


✓125.65
✓126.98
✓134.94

✓149.53
✓152.27


Parameter	Value
1 Title	SPK01253P3.fid
2 Solvent	CDCl ₃
3 Temperature	298.0
4 Number of Scans	1024
5 Receiver Gain	2050.0
6 Relaxation Delay	5.0000
7 Pulse Width	10.6300
8 Spectrometer Frequency	150.97
9 Nucleus	¹³ C





Parameter	Value
1 Title	SBK01253P2.fid
2 Solvent	CDCl ₃
3 Temperature	298.0
4 Number of Scans	128
5 Receiver Gain	362.0
6 Relaxation Delay	1.0000
7 Pulse Width	11.0000
8 Spectrometer Frequency	192.61
9 Nucleus	11B



—30.60



Parameter	Value
1 Title	SBK01256P1.fid
2 Solvent	CDCl ₃
3 Temperature	298.0
4 Number of Scans	16
5 Receiver Gain	144.0
6 Relaxation Delay	1.0000
7 Pulse Width	10.7700
8 Spectrometer Frequency	600.32
9 Nucleus	¹ H





✓ 24.97
✓ 21.31
✓ 19.88


—83.37

✓ 147.14
✓ 138.67
✓ 136.36
✓ 134.02
✓ 131.32
✓ 127.03
✓ 125.88


Parameter	Value
1 Title	SBK01256P.3.fid
2 Solvent	CDCl ₃
3 Temperature	298.0
4 Number of Scans	1024
5 Receiver Gain	2050.0
6 Relaxation Delay	5.0000
7 Pulse Width	10.6300
8 Spectrometer Frequency	150.97
9 Nucleus	¹³ C





Parameter	Value
1 Title	SBK01265-boron.1.fid
2 Solvent	CDCl ₃
3 Temperature	299.4
4 Number of Scans	128
5 Receiver Gain	203.0
6 Relaxation Delay	1.0000
7 Pulse Width	8.3000
8 Spectrometer Frequency	128.39
9 Nucleus	11B




—29.61





Parameter	Value
1 Title	SBK01260B.1.fid
2 Solvent	CDCl ₃
3 Temperature	298.0
4 Number of Scans	16
5 Receiver Gain	161.0
6 Relaxation Delay	1.0000
7 Pulse Width	10.7700
8 Spectrometer Frequency	600.32
9 Nucleus	¹ H







-30.31

Parameter	Value
1 Title	SBK01260B.2.fid
2 Solvent	DCI3
3 Temperature	298.0
4 Number of Scans	128
5 Receiver Gain	322.0
6 Relaxation Delay	1.0000
7 Pulse Width	11.0000
8 Spectrometer Frequency	192.61
9 Nucleus	11B












-29.77

Parameter	Value
1 Title	SBK01276-boron.1.fid
2 Solvent	CDCl ₃
3 Temperature	299.4
4 Number of Scans	128
5 Receiver Gain	203.0
6 Relaxation Delay	1.0000
7 Pulse Width	8.3000
8 Spectrometer Frequency	128.39
9 Nucleus	11B













Parameter	Value
1 Title	SBK01274.10 fid
2 Solvent	CDCl ₃
3 Temperature	297.4
4 Number of Scans	16
5 Receiver Gain	181.0
6 Relaxation Delay	1.0000
7 Pulse Width	10.7700
8 Spectrometer Frequency	600.32
9 Nucleus	¹ H








Parameter	Value
1 Title	SBK01274.11.fid
2 Solvent	CDCl ₃
3 Temperature	297.4
4 Number of Scans	128
5 Receiver Gain	362.0
6 Relaxation Delay	1.0000
7 Pulse Width	11.0000
8 Spectrometer Frequency	192.61
9 Nucleus	11B




Parameter	Value
1 Title	SBK01283A-F2.1.fid
2 Solvent	CDCl ₃
3 Temperature	298.0
4 Number of Scans	16
5 Receiver Gain	181.0
6 Relaxation Delay	1.0000
7 Pulse Width	10.7700
8 Spectrometer Frequency	600.32
9 Nucleus	¹ H







Parameter	Value
1 Title	SBK01283-600.2.fid
2 Solvent	CDCl ₃
3 Temperature	297.6
4 Number of Scans	128
5 Receiver Gain	322.0
6 Relaxation Delay	1.0000
7 Pulse Width	11.0000
8 Spectrometer Frequency	192.61
9 Nucleus	11B




-30.27





Parameter	Value
1 Title	SBK01283-400.3.fid
2 Solvent	CDCl ₃
3 Temperature	298.1
4 Number of Scans	16
5 Receiver Gain	1625.5
6 Relaxation Delay	2.0000
7 Pulse Width	15.0300
8 Spectrometer Frequency	376.46
9 Nucleus	¹⁹ F




—112.37









Parameter	Value
1 Title	WBR03149F2again.2.fid
2 Solvent	CDCl ₃
3 Temperature	297.1
4 Number of Scans	128
5 Receiver Gain	362.0
6 Relaxation Delay	1.0000
7 Pulse Width	11.0000
8 Spectrometer Frequency	192.61
9 Nucleus	11B



—30.24







145.31
140.43
135.76
135.03
130.09
128.85
126.37
125.17
123.86
121.59
120.95
113.84


—24.97
—21.74

—83.55


Parameter	Value
1 Title	SBK01255.1.fid
2 Solvent	CDCl ₃
3 Temperature	298.2
4 Number of Scans	1024
5 Receiver Gain	512.0
6 Relaxation Delay	3.0000
7 Pulse Width	9.2500
8 Spectrometer Frequency	100.62
9 Nucleus	¹³ C



Parameter	Value
1 Title	SBK01255-boron.1.fid
2 Solvent	CHCl ₃
3 Temperature	299.4
4 Number of Scans	128
5 Receiver Gain	203.0
6 Relaxation Delay	1.0000
7 Pulse Width	8.3000
8 Spectrometer Frequency	128.39
9 Nucleus	11B



-29.15





7.03
7.04
7.05
7.06
7.07
7.08
7.09
7.10
7.11
7.12
7.13
7.14
7.15
7.16
7.17
7.18
7.19
7.20
7.21
7.22
7.23
7.24
7.25
7.26
7.27
7.28
7.29
7.30
7.31
7.32
7.33
7.34
7.35
7.36
7.37
7.38
7.39
7.40
7.41
7.42
7.43
7.44
7.45
7.46
7.47
7.48
7.49
7.50
7.51
7.52
7.53
7.54
7.55
7.56
7.57
7.58
7.59
7.60
7.61
7.62
7.63
7.64
7.65
7.66
7.67
7.68
7.69
7.70
7.71
7.72
7.73
7.74
7.75
7.76
7.77
7.78
7.79
7.80
7.81
7.82
7.83
7.84
7.85
7.86
7.87
7.88
7.89
7.90
7.91
7.92
7.93
7.94
7.95
7.96
7.97
7.98
7.99
8.00
8.01
8.02
8.03
8.04
8.05
8.06
8.07
8.08
8.09
8.10
8.11
8.12
8.13
8.14
8.15
8.16
8.17
8.18
8.19
8.20
8.21
8.22
8.23
8.24
8.25
8.26
8.27
8.28
8.29
8.30
8.31
8.32
8.33
8.34
8.35
8.36
8.37
8.38
8.39
8.40
8.41
8.42
8.43
8.44
8.45
8.46
8.47
8.48
8.49
8.50
8.51
8.52
8.53
8.54
8.55
8.56
8.57
8.58
8.59
8.60
8.61
8.62
8.63
8.64
8.65
8.66
8.67
8.68
8.69
8.70
8.71
8.72
8.73
8.74
8.75
8.76
8.77
8.78
8.79
8.80
8.81
8.82
8.83
8.84
8.85
8.86
8.87
8.88
8.89
8.90
8.91
8.92
8.93
8.94
8.95
8.96
8.97
8.98
8.99
9.00
9.01
9.02
9.03
9.04
9.05
9.06
9.07
9.08
9.09
9.10
9.11
9.12
9.13
9.14
9.15
9.16
9.17
9.18
9.19
9.20
9.21
9.22
9.23
9.24
9.25
9.26
9.27
9.28
9.29
9.30
9.31
9.32
9.33
9.34
9.35
9.36
9.37
9.38
9.39
9.40
9.41
9.42
9.43
9.44
9.45
9.46
9.47
9.48
9.49
9.50
9.51
9.52
9.53
9.54
9.55
9.56
9.57
9.58
9.59
9.60
9.61
9.62
9.63
9.64
9.65
9.66
9.67
9.68
9.69
9.70
9.71
9.72
9.73
9.74
9.75
9.76
9.77
9.78
9.79
9.80
9.81
9.82
9.83
9.84
9.85
9.86
9.87
9.88
9.89
9.90
9.91
9.92
9.93
9.94
9.95
9.96
9.97
9.98
9.99
10.00

Parameter	Value
1 Title	SBK01254P.1.fid
2 Solvent	CDCl ₃
3 Temperature	298.0
4 Number of Scans	16
5 Receiver Gain	181.0
6 Relaxation Delay	1.0000
7 Pulse Width	10.7700
8 Spectrometer Frequency	600.32
9 Nucleus	¹ H




-1.31








Parameter	Value
1 Title	SBK01254P2.fid
2 Solvent	CDCl ₃
3 Temperature	298.0
4 Number of Scans	128
5 Receiver Gain	322.0
6 Relaxation Delay	1.0000
7 Pulse Width	11.0000
8 Spectrometer Frequency	192.61
9 Nucleus	11B




—30.37







Parameter	Value
1 Title	SBK01264-F1-2.10.fid
2 Solvent	CDCl3
3 Temperature	298.0
4 Number of Scans	16
5 Receiver Gain	32.0
6 Relaxation Delay	10.0000
7 Pulse Width	10.7700
8 Spectrometer Frequency	600.32
9 Nucleus	1H




Parameter	Value
1 Title	SBK01264-F1-2.11.fid
2 Solvent	CDCl ₃
3 Temperature	298.0
4 Number of Scans	128
5 Receiver Gain	362.0
6 Relaxation Delay	1.0000
7 Pulse Width	11.0000
8 Spectrometer Frequency	192.61
9 Nucleus	11B





—30.04







Parameter	Value
1 Title	SBK01301-13C.2.fid
2 Solvent	CDCB
3 Temperature	301.2
4 Number of Scans	32
5 Receiver Gain	512.0
6 Relaxation Delay	3.0000
7 Pulse Width	9.2500
8 Spectrometer Frequency	100.62
9 Nucleus	13C




Parameter	Value
1 Title	SBK01301-boron.1.fid
2 Solvent	CDCl ₃
3 Temperature	299.4
4 Number of Scans	128
5 Receiver Gain	203.0
6 Relaxation Delay	1.0000
7 Pulse Width	8.3000
8 Spectrometer Frequency	128.39
9 Nucleus	11B





-25.63







Parameter	Value
1 Title	SBK01285.3.fid
2 Solvent	CDCl ₃
3 Temperature	299.0
4 Number of Scans	1024
5 Receiver Gain	2050.0
6 Relaxation Delay	5.0000
7 Pulse Width	10.6300
8 Spectrometer Frequency	150.97
9 Nucleus	¹³ C





Parameter	Value
1 Title	SBK01285.2.fid
2 Solvent	CDCl ₃
3 Temperature	297.3
4 Number of Scans	128
5 Receiver Gain	362.0
6 Relaxation Delay	1.0000
7 Pulse Width	11.0000
8 Spectrometer Frequency	192.61
9 Nucleus	11B




—26.26





Parameter	Value
1 Title	SBK01293.1.fid
2 Solvent	CDCl ₃
3 Temperature	298.0
4 Number of Scans	16
5 Receiver Gain	181.0
6 Relaxation Delay	1.0000
7 Pulse Width	10.7700
8 Spectrometer Frequency	600.32
9 Nucleus	¹ H







-30.80


Parameter	Value
1 Title	SBK01293.2.fid
2 Solvent	DCIβ
3 Temperature	298.0
4 Number of Scans	128
5 Receiver Gain	362.0
6 Relaxation Delay	1.0000
7 Pulse Width	11.0000
8 Spectrometer Frequency	192.61
9 Nucleus	11B




—7.47
—7.45
—7.40
—7.38
—7.36
—7.34
—7.17
—7.14
—7.12
—7.10
—7.03
—7.01
—6.38
—6.36
—6.31
—6.26
—5.86

Parameter	Value
1 Title	SBK01297.1.fid
2 Solvent	CDCl ₃
3 Temperature	298.2
4 Number of Scans	16
5 Receiver Gain	9.0
6 Relaxation Delay	1.0000
7 Pulse Width	15.0000
8 Spectrometer Frequency	400.13
9 Nucleus	¹ H






Parameter	Value
1 Title	SBK01297-11B.1.fid
2 Solvent	CDCl ₃
3 Temperature	299.0
4 Number of Scans	128
5 Receiver Gain	322.0
6 Relaxation Delay	1.0000
7 Pulse Width	11.0000
8 Spectrometer Frequency	192.61
9 Nucleus	11B




—28.88



6.63
6.30
6.27


—1.28

Parameter	Value
1 Title	SBK01298.1.fid
2 Solvent	Acetone
3 Temperature	298.0
4 Number of Scans	16
5 Receiver Gain	181.0
6 Relaxation Delay	1.0000
7 Pulse Width	10.7700
8 Spectrometer Frequency	600.32
9 Nucleus	¹ H




149.11
139.16
134.32
134.28
134.23
134.18
126.17
125.71

Parameter	Value
1 Title	SBK01298-13C.1.fid
2 Solvent	Acetone
3 Temperature	301.2
4 Number of Scans	1024
5 Receiver Gain	512.0
6 Relaxation Delay	3.0000
7 Pulse Width	9.2500
8 Spectrometer Frequency	100.62
9 Nucleus	¹³ C



-2.98



Parameter	Value
1 Title	SBK01298.2.fid
2 Solvent	Acetone
3 Temperature	298.0
4 Number of Scans	128
5 Receiver Gain	362.0
6 Relaxation Delay	1.0000
7 Pulse Width	11.0000
8 Spectrometer Frequency	192.61
9 Nucleus	1H



Parameter	Value
1 Title	SBK01298-19F.1.fid
2 Solvent	Acetone
3 Temperature	301.2
4 Number of Scans	16
5 Receiver Gain	1625.5
6 Relaxation Delay	2.0000
7 Pulse Width	15.0300
8 Spectrometer Frequency	376.46
9 Nucleus	¹⁹ F



—141.57

