

Rapid isolation of dengue-neutralizing antibodies from single cell-sorted human antigen-specific memory B-cell cultures

Kara S. Cox^a, Aimin Tang^a, Zhifeng Chen^a, Melanie S. Horton^a, Hao Yan^a, Xin-Min Wang^a, Sheri A. Dubey^a, Daniel J. DiStefano^a, Andrew Ettenger^b, Rachel H. Fong^b, Benjamin J. Doranz^b, Danilo R. Casimiro^a and Kalpit A. Vora^a

^a Department of Infectious Diseases and Vaccines-West Point, PA, Merck Research Laboratories, Merck & Co., Inc. Kenilworth, NJ USA

^b Integral Molecular Inc. , Philadelphia, PA USA

Corresponding author: Kalpit A. Vora; kalpit.vora@merck.com

Supplemental Table 1

Mab	HCDR3	Kappa CDR3	Lambda CDR3
DB5	DDGGSYGFDY		QTWDTTTHVV
DB6	PITSVAGGGTWFD P	QQYDSFPIT	
DC11	QEVAVAGNSY	QQRNYWPHT	
DC4	RPWTMPSDY	QQYDNWPWT	
DC6	RPWTYAHFDY	QQYGSSPLT	
DH1	DCSPTTCYFQ	QQSYNDPRT	
DD1L	GEQWPSDF		ATWDDRVSGPV
DD1K	DSHQWLAYDF	QQRSNWPRDT	
DD12	TSNRDGPLVDY	QQTYITPRT	

Table S1 Monoclonal Antibody CDR_{3H} and CDR_{3L} sequences

Heavy and Light chain sequences from the CDR3 regions of each of the human mAbs which displayed neutralizing activity are listed.