MICROBIOLOGY, BIOCHEMISTRY, STRUCTURAL BIOLOGY

*Hyun Ho Park, Phone: +82-53-810-3045. Fax: +82-53-810-4516. E-mail: hyunho@ynu.ac.kr *Tae-Jin Lee, Phone: +82-53-640-6914. Fax:+82-53-621-5053. E-mail: tjlee@med.yu.ac.kr

Rescuing neuronal cell death by RAIDD- and PIDD- derived peptides and its implications for therapeutic intervention in neurodegenerative diseases

Tae-Ho Jang $^{a,\#}$, In-Hye Lim $^{b,\#}$, Chang Min Kim $^{a,\#}$, Jae Young Choi a , Eun-Ae Kim b , Tae-Jin Lee b,* and Hyun Ho Park a,*

^aSchool of Biotechnology and Graduate School of Biochemistry at Yeungnam University, Gyeongsan 38541, South Korea

^bDepartment of Anatomy, College of Medicine, Yeungnam University, 317-1 Daemyung-Dong Nam-Gu, Daegu 42415, South Korea




Figure S1. Peptide (PDH3) dose-dependent inhibition of PIDDosome core formation at low concentration range. The RAIDD DD:PIDD DD complex was prepared and mixed with the concentrations of PDH3 peptide indicated above the gel, after which the mixtures were subjected to native-PAGE.




Figure S2. Peptides specificity analysis. Another DD family, RIP1 DD, FADD DD, was prepared for the peptides specificity analysis with Native-PAGE.