

1 **Gene transduction in mammalian cells using *Bombyx mori***
2 **nucleopolyhedrovirus assisted by glycoprotein 64 of**
3 ***Autographa californica* multiple nucleopolyhedrovirus**

4
5 Tatsuya Kato,^{1,2,3*} Saki Sugioka,² Kohei Itagaki,² and Enoch Y. Park^{1,2,3*}

6
7 ¹ Laboratory of Biotechnology, Green Chemistry Research Division, Research
8 Institute of Green Science and Technology, Shizuoka University, 836 Ohya
9 Suruga-ku, Shizuoka, 422-8529, Japan

10 ² Laboratory of Biotechnology, Department of Applied Biological Chemistry,
11 Faculty of Agriculture, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka, 422-
12 8529, Japan

13 ³ Laboratory of Biotechnology, Department of Bioscience, Graduate School of
14 Science and Technology, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka,
15 422-8529, Japan

16
17 E-mail:

18 kato.tatsuya@shizuoka.ac.jp (TK)

19 sakicsxxx@yahoo.co.jp (SS)


20 pixy_007bc@yahoo.co.jp (KI)

21 park.enoch@shizuoka.ac.jp (EYP)

22

* Correspondence

Tatsuya Kato (kato.tatsuya@shizuoka.ac.jp); Enoch Y. Park (park.enoch@shizuoka.ac.jp)



24

25

26 **Figure S1. Fluorescence microscopy of HEK293T cells transduced with**
 27 **BmNPV Δ bgrp/AcGP64/EGFP.** Transduction of recombinant
 28 BmNPV Δ bgrp/AcGP64/EGFPs into HEK293T cells at an M.O.I. of 300. Each
 29 transduction efficiency of recombinant BmNPV Δ bgrp/AcGP64/EGFPs into HEK293T
 30 cells was 7.8%, 9.5%, 12.7% and 8.3% judging by counting the number of fluorescent
 31 cells.