

## AR relation with diabetes in TNBC

**Supplementary Table 1.** Clinical pathological features of TNBC samples

	Patients (n=238)
<b>Age</b>	
<40	26 (10.9%)
≥40≤60	107 (45.0%)
>60	101 (42.4%)
Unknown	4 (1.7%)
<b>Grade</b>	
I-II	28 (11.8%)
III	197 (82.8%)
Unknown	13 (5.5%)
<b>Size (T)</b>	
≤2 cm	102 (42.9%)
>2≤5 cm	103 (43.3%)
>5	20 (8.4%)
Unknown	13 (5.5%)
<b>LNM</b>	
Yes	81 (34.0%)
No	111 (46.6%)
Unknown	46 (19.3%)
<b>Metastasis</b>	
Yes	42 (17.6%)
No	131 (55.0%)
Unknown	65 (27.3%)
<b>Ki-67</b>	
>20	186 (78.2%)
≤20	42 (17.6%)
Unknown	10 (4.2%)
<b>BMI</b>	
<30	72 (30.3%)
≥30	36 (15.1%)
Unknown	130 (54.6%)
<b>Diabetic</b>	
Yes	17 (7.1%)
No	98 (41.2%)
Unknown	123 (51.7%)
<b>Follow up</b>	
NED	63 (26.5%)
AWD	24 (10.1%)
DOD	16 (6.7%)
Unknown	135 (56.7%)

LNM=Lymph node metastasis; NED=No evidence of disease; AWD=Alive with disease; DOD=Dead of disease.