





**A****B**

**Supplementary Data Figure 1. The mouse cell line GL261 and RCAS-PDGFB murine tumor model used in the present study contains a population of CD133<sup>+</sup> cells with increased tumorigenic potential, which is lost after differentiation.**


(A) The percentage of CD133<sup>+</sup> cells in neurosphere GL261 cells is 30,1% while in adhesive GL261 it is less than 1%. The mouse astrocytoma cell line GL261 contained a population of CD133<sup>+</sup> cells, which could be purified by FACS with respective species-specific antibodies. The respective immunoglobulin isotypes (isotype control) served as controls. (B) RCAS-PDGFB murine tumor model contained a population of CD133<sup>+</sup> cells, which could be purified by FACS with respective species-specific antibodies. The respective immunoglobulin isotypes (isotype control) served as controls.



**Supplementary Data Figure 2. Flow cytometric characterization of FACS sorted CD133<sup>+</sup> GL261 cells**  
 Representative histograms of GFAP, Sox2, Notch1 and CD133 expression of FACS sorted CD133<sup>+</sup> GL261 cultured in stem cell medium (non-differentiated) and differentiated with 10% FCS. The gray label indicates isotype control. These data indicate that CD133 positive cells have the capacity to differentiate.



**Supplementary Data Figure 3. Protein concentrations of the conditioned media are equal**  
The protein concentration of the conditioned media were determined using the BCA protein assay reagent from Pierce.



**Supplementary Data Figure 4. Expression levels of selected TLR4-specific ligands in GL261 CD133<sup>+</sup> cells.**

Expression of Fibronectin EDA-domain, Hsp22 and tenascin-C was analyzed by qRT-PCR in FACS sorted CD133<sup>-</sup> and CD133<sup>+</sup> cells from mouse GL261 cell cultures.  $\beta$ -Actin was used as an internal control, target expression in CD133<sup>+</sup> cells was normalized to the respective CD133<sup>-</sup> fraction. Graph represents the fold change from n=7 ( $\pm$ SD) individual CD133<sup>+</sup> cells compared to the CD133<sup>-</sup> fraction isolated from the same sample respectively. HSP22 and EDA are slightly higher expressed in CD133<sup>+</sup> compared to CD133<sup>-</sup> cells, while tenascin-C is significant 4 fold higher expressed in CD133<sup>+</sup> compared to CD133<sup>-</sup> cells.

<b>Gender</b>	<b>Age</b>	<b>Diagnose</b>	<b>Code in text</b>
Female	47	GBM	GBM 1
Female	62	GBM	GBM 2
Male	78	GBM	GBM 3
Female	71	GBM	GBM 4
Female	54	GBM	GBM 5
Male	54	GBM	GBM 6
Female	74	GBM	GBM 7
Female	31	GBM	GBM 8

**Supplementary Data Table 1. Overview of resected tumor tissue samples used for qRT-PCR**