Elastic instabilities in planar elongational flow of monodisperse polymer solutions

Simon J. Haward^{1,*}, Gareth H. McKinley², and Amy Q. Shen¹

Supplementary Information List of Video Captions

Movie 1 (M1.mov): Time-resolved flow velocimetry for flow of a 0.07 wt% solution of PS16 in DOP at Wi = 0.39, Re = 0.019, and corresponding to the space-time data shown in Fig. 8(a) of the main article.

Movie 2 (M2.mov): Time-resolved flow velocimetry for flow of a 0.07 wt% solution of PS16 in DOP at Wi = 1.17, Re = 0.058, and corresponding to the space-time data shown in Fig. 8(b) of the main article.

Movie 3 (M3.mov): Time-resolved flow velocimetry for flow of a 0.07 wt% solution of PS16 in DOP at Wi = 1.96, Re = 0.10, and corresponding to the space-time data shown in Fig. 8(c) of the main article.

Movie 4 (M4.mov): Time-resolved flow velocimetry for flow of a 0.07 wt% solution of PS16 in DOP at Wi = 2.74, Re = 0.14, and corresponding to the space-time data shown in Fig. 8(d) of the main article.

¹Okinawa Institute of Science and Technology, Onna-son, Okinawa, 904-0495, Japan ²Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, United States *simon.haward@oist.jp