

Gene	NM	Primer sequences (Fwd./Rev.)	Expression in the liver
<b>Acta2</b>	007392.3	CCAGCACCATGAAGATCAAG TGGAAGGTAGACAGCGAAGC	Enhanced expression in activated myofibroblasts/HSCs[2]
<b>CK19</b>	008471.2	ATGAGATCATGGCCGAGAAG GGTGTTCAGCTCCTCAATCC	Marker of biliary epithelial cells and liver progenitor cells[3]
<b>F4/80</b>	010130.4	CCTGGACGAATCCTGTGAAG GGTGGGACCACAGAGAGTTG	Marker of hepatic macrophages[4]
<b>Colla1</b>	007742.4	ACCTAAGGGTACCGCTGGA TCCAGCTTCTCCATCTTTGC	Enhanced expression in activated myofibroblasts/HSCs[2]
<b>Gapdh</b>	008084.2	CCTGCTTCACCACCTTCTTG TGTCCTCGTGGATCTGAC	Used as reference gene
<b>Lamc1</b>	010683.2	TTTGATAGACGCGTGAACGATAA TGGCGGAATTCTCCTTAGA	Enhanced in fibrotic livers[5, 6]
<b>Cyp3a11</b>	007818.3	TGAATATGAAACTTGCTCTCACTAAAA CCTTGTCTGCTTAATTCAGAGGT	Marker of hepatocytes[7]
<b>Stab2</b>	138673.2	AGCTGCTGCCTTTAATCCTCA ACTCCGTCTTGATGGTTAGAGTA	Markers for liver sinusoidal cells[8, 9]
<b>Epcam</b>	008532.2	GCGGCTCAGAGAGACTGTG CCAAGCATTAGACGCCAGTTT	Marker of biliary epithelial cells and liver progenitor cells[10]

### Supplementary Table S2: Oligonucleotides used for RT-qPCR.

- Mannaerts I, Leite SB, Verhulst S, Claerhout S, Eysackers N, Thoen LF, et al. The Hippo pathway effector YAP controls mouse hepatic stellate cell activation. *J Hepatol.* 2015;63(3):679-88. doi: 10.1016/j.jhep.2015.04.011. PubMed PMID: 25908270.
- Yimlamai D, Christodoulou C, Galli GG, Yanger K, Pepe-Mooney B, Gurung B, et al. Hippo pathway activity influences liver cell fate. *Cell.* 2014;157(6):1324-38. doi: 10.1016/j.cell.2014.03.060. PubMed PMID: 24906150; PubMed Central PMCID: PMC4136468.
- Lee SH, Starkey PM, Gordon S. Quantitative-Analysis of Total Macrophage Content in Adult-Mouse Tissues - Immunochemical Studies with Monoclonal-Antibody F4/80. *Journal of Experimental Medicine.* 1985;161(3):475-89. doi: DOI 10.1084/jem.161.3.475. PubMed PMID: WOS:A1985ACS5200004.
- Kallis YN, Robson AJ, Fallowfield JA, Thomas HC, Alison MR, Wright NA, et al. Remodelling of extracellular matrix is a requirement for the hepatic progenitor cell response. *Gut.* 2011;60(4):525-33. doi: 10.1136/gut.2010.224436. PubMed PMID: 21106552.
- Knittel T, Janneck T, Muller L, Fellmer P, Ramadori G. Transforming growth factor beta 1-regulated gene expression of Ito cells. *Hepatology.* 1996;24(2):352-60. doi: 10.1053/jhep.1996.v24.pm0008690404. PubMed PMID: 8690404.
- Anzai K, Chikada H, Tsuruya K, Ida K, Kagawa T, Inagaki Y, et al. Foetal hepatic progenitor cells assume a cholangiocytic cell phenotype during two-dimensional pre-culture. *Sci Rep.* 2016;6:28283. doi: 10.1038/srep28283. PubMed PMID: 27335264; PubMed Central PMCID: PMC4917868.
- Hansen B, Longati P, Elvevold K, Nedredal GI, Schledzewski K, Olsen R, et al. Stabilin-1 and stabilin-2 are both directed into the early endocytic pathway in hepatic sinusoidal endothelium via interactions with clathrin/AP-2, independent of ligand binding. *Exp Cell Res.* 2005;303(1):160-73. doi: 10.1016/j.yexcr.2004.09.017. PubMed PMID: 15572036.
- Nonaka H, Sugano S, Miyajima A. Serial analysis of gene expression in sinusoidal endothelial cells from normal and injured mouse liver. *Biochem Biophys Res Commun.* 2004;324(1):15-24. doi: 10.1016/j.bbrc.2004.09.014. PubMed PMID: 15464976.
- Lu WY, Bird TG, Boulter L, Tsuchiya A, Cole AM, Hay T, et al. Hepatic progenitor cells of biliary origin with liver repopulation capacity. *Nat Cell Biol.* 2015;17(8):971-83. doi: 10.1038/ncb3203. PubMed PMID: 26192438; PubMed Central PMCID: PMC4612439.