

eTable 3. Sex- and age-specific PAR (%) associated with selected cardiovascular risk factors for fatal CHD and stroke in the Indonesian population

	Smoking	Hypertension	Elevated total cholesterol	Excess body weight	Diabetes
Men					
Fatal CHD	28.0 (20.1, 35.4)	20.1 (16.0, 23.8)	13.0 (7.4, 18.5)	7.7 (4.0, 11.8)	6.4 (3.7, 10.1)
Fatal Stroke	17.2 (12.0, 22.5)	40.1 (37.4, 42.8)	3.5 (-0.2, 7.1)	2.5 (0.1, 4.9)	4.2 (2.4, 6.4)
Women					
Fatal CHD	1.3 (0.8, 1.7)	24.1 (19.8, 28.5)	16.7 (9.2, 23.2)	12.1 (6.4, 18.3)	12.0 (5.5, 20.3)
Fatal Stroke	0.7 (0.4, 0.9)	45.8 (42.6, 48.6)	4.4 (0.3, 8.9)	4.0 (0.3, 8.0)	7.2 (3.2, 12.2)
<55 years					
Fatal CHD	22.7 (12.8, 33.5)	36.2 (29.9, 42.4)	17.9 (9.7, 27.1)	17.4 (8.7, 27.0)	11.0 (6.5, 16.7)
Fatal Stroke	13.1 (6.2, 19.6)	59.1 (55.4, 62.7)	7.2 (1.4, 13.4)	10.2 (4.1, 16.7)	8.3 (5.0, 12.1)
≥55 years					
Fatal CHD	8.4 (2.5, 14.5)	33.1 (25.4, 40.1)	15.7 (4.5, 15.6)	3.7 (-0.7, 8.6)	12.5 (6.7, 19.4)
Fatal Stroke	2.9 (-0.5, 6.4)	55.4 (51.7, 58.9)	2.7 (-4.3, 9.3)	-1.4 (-4.1, 1.5)	6.0 (2.2, 10.3)

CHD, coronary heart disease; PAR, population attributable risk.