

Journal name: PLANTA

Article title: Immunolocalization of cyclotides in plant cells, tissues and organs supports their role in host defense

Authors: Blazej Slazak^{1,2}, Małgorzata Kapusta³, Sohaib Malik², Jerzy Bohdanowicz³, Elżbieta Kuta⁴, Przemysław Malec⁵, Ulf Göransson²

Affiliations:

¹W. Szafer Institute of Botany, Polish Academy of Science, 46 Lubicz St., 31-512 Cracow, Poland

²Division of Pharmacognosy, Department of Medicinal Chemistry, Uppsala University, Biomedical Center, Box 574, 751 23 Uppsala, Sweden

³Department of Plant Cytology and Embryology, Faculty of Biology, University of Gdańsk, 59 Wita Stwosza St., 80-308 Gdańsk, Poland

⁴Department of Plant Cytology and Embryology, Institute of Botany, Jagiellonian University, 9 Gronostajowa St., 30-387 Cracow, Poland

⁵Faculty of Biochemistry, Biophysics and Biotechnology; Jagiellonian University, 7 Gronostajowa St, 30-387 Cracow, Poland

Corresponding author: Blazej Slazak

¹W. Szafer Institute of Botany, Polish Academy of Science, 46 Lubicz St., 31-512 Cracow, Poland

phone: +48 660 450 799

e-mail: b.slazak@botany.com

Supplement 1. Additional dot and western blot negative controls confirming the antibody's specificity. **a**, no signal (violet color reaction) can be detected on the membrane with blotted 2 μ g of ovalbumin and BSA. **b, c**, western blots stained with Ponceau S for whole blotted protein (**b**) and using the anti-cyO2 primary antibody and a secondary antibody conjugated with ALP (violet color reaction marked with arrowheads)(**c**). Lanes 1, 2, 3 and 4 correspond to *A. thaliana*, *N. benthamiana*, *V. uliginosa*, and pure cyO2, respectively.

