

SUPPLEMENTARY TABLE S2. THE RATIO (MEDIAN%) OF LYMPHOCYTE SUBSETS TO THE PARENT CELL POPULATION AMONG PWID AND CONTROLS

<i>Cell populations</i>	<i>Triple negative PWID (n=11)</i>	<i>Controls</i>	<i>p</i>
CD4 ⁺			
T _N	58.6 (56.6–64.9)	55.9 (49.5–63.3)	0.150
T _N HLA-DR ⁺	36.4 (33.5–52.9)	32.2 (24.4–39.0)	0.128
T _N CCR5 ⁺	1.5 (1.0–1.9)	1.7 (1.0–2.3)	0.445
T _{CM}	0.3 (0.1–1.1)	0.3 (0.2–0.5)	0.606
T _{CM} HLA-DR ⁺	44.0 (37.0–53.9)	48.9 (40.8–54.5)	0.625
T _{CM} CCR5 ⁺	4.9 (3.7–6.0)	3.5 (2.7–4.0)	0.020
T _{EM}	2.3 (1.3–2.6)	1.3 (0.6–2.0)	0.020
T _{EM} HLA-DR ⁺	10.8 (8.9–15.4)	16.4 (11.7–21.9)	0.031
T _{EM} CCR5 ⁺	23.2 (21.0–25.9)	15.0 (11.3–18.8)	<0.001
T _{TEMRA}	10.9 (8.2–12.6)	8.0 (3.7–10.9)	0.044
T _{TEMRA} HLA-DR ⁺	0.5 (0.1–2.2)	0.3 (0.2–0.8)	0.625
T _{TEMRA} CCR5 ⁺	22.2 (16.9–37.5)	11.1 (3.9–20.7)	0.016
HLA-DR ⁺	1.5 (0–8.7)	0 (0–5.4)	0.102
CCRS ⁺	5.4 (4.1–8.7)	4.6 (3.8–5.4)	0.222
CCRS ⁺	2.4 (1.6–3.4)	2.0 (1.0–2.9)	0.307
CD8 ⁺			
T _N	31.5 (25.6–34.1)	32.3 (28.2–41.2)	0.167
T _N HLA-DR ⁺	25.8 (16.8–39.2)	23.2 (13.5–35.7)	0.639
T _N CCR5 ⁺	2.8 (1.1–3.8)	2.4 (1.6–3.1)	0.653
T _{CM}	0.7 (0.1–1.0)	0.3 (0.1–0.6)	0.292
T _{CM} HLA-DR ⁺	3.9 (3.1–5.4)	2.4 (1.6–3.7)	0.017
T _{CM} CCR5 ⁺	9.5 (5.3–12.9)	7.2 (4.1–10.0)	0.219
T _{EM}	3.2 (2.3–8.7)	1.8 (0–5.6)	0.036
T _{EM} HLA-DR ⁺	31.6 (27.4–40.3)	43.2 (30.0–54.0)	0.118
T _{EM} CCR5 ⁺	31.3 (23.5–38.6)	18.6 (11.7–24.5)	0.002
T _{TEMRA}	16.4 (14.5–24.0)	15.8 (9.2–24.9)	0.399
T _{TEMRA} HLA-DR ⁺	29.2 (24.3–35.9)	27.1 (16.8–35.6)	0.367
T _{TEMRA} CCR5 ⁺	30.8 (18.9–42.3)	13.0 (7.6–19.5)	<0.001
HLA-DR ⁺	4.4 (3.4–9.5)	3.3 (1.2–6.1)	0.161
CCRS ⁺	23.9 (15.2–27.9)	10.5 (8.1–15.9)	0.002
CCRS ⁺	8.7 (6.4–13.2)	8.1 (4.1–11.2)	0.421

p<0.05 was considered statistically significant and is marked in bold typeface.