

Table 1. Select demographic and clinical characteristics of sampled patients, sero-survey of *Cryptococcus* antigenemia among HIV-infected adults with advanced immunosuppression in Namibia, 2013-14.

Variable	Result
Patient samples included, #	825
Sex, #, [% (95% CI)]^a	
Female	374, [45.9 (42.5 – 49.4)]
Male	440, [54.1 (50.6 – 57.5)]
Age, median (min - max) (IQR) years^b	38 (15 - 94) (32 - 46)
Age group, #, [% (95% CI)]	
15 - 24 years	50 [6.2 (4.7 – 8.0)]
25 - 34 years	224 [27.6 (24.6 – 30.7)]
≥ 35 years	539 [66.3 (63.0 – 69.5)]
CD4⁺ result, median (IQR) cells/µL^c	85 (51 - 114)
CD4⁺ result strata 1, #, [% (95% CI)]	
< 100 cells/µL	511, [62.9 (59.7 -66.4)]
100 - 200 cells/µL	302, [37.1 (33.9 – 40.5)]
CD4⁺ result strata 2, #, [% (95% CI)]	
< 50 cells/µL	194, [23.5 (20.7 – 26.6)]
50 - 200 cells/µL	630, [76.5 (73.4 – 79.3)]

^a. data missing, n=11

^b. data missing, n=12

^c. data missing, n=12