Supplemental Information

Suppression of iASPP-dependent aggressiveness in cervical cancer through reversal of methylation silencing of microRNA-124

Peixin Dong^{1, *}, Ying Xiong^{2, *}, Hidemichi Watari³, Sharon JB Hanley¹, Yosuke Konno³, Kei Ihira³, Fumihiko Suzuki⁴, Takahiro Yamada¹, Masataka Kudo³, Junming Yue⁵ & Noriaki Sakuragi^{1, 3}

¹Department of Women's Health Educational System, Hokkaido University School of Medicine, Hokkaido University, N15, W7, Sapporo 0608638, Japan

²Department of Gynecology, State Key Laboratory of Oncology in South China, Sun Yat-sen University Cancer Center, Guangzhou 510060, P. R. China


³Department of Gynecology, Hokkaido University School of Medicine, Hokkaido University, N15, W7, Sapporo 0608638, Japan

⁴Department of Obstetrics and Gynecology, Tohoku University, Sendai 9808574, Japan


⁵Department of Pathology and Laboratory Medicine, University of Tennessee Health Science Center, USA; Center for Cancer Research, University of Tennessee Health Science Center, Memphis, TN, USA

*Equal contributors

Correspondence should be addressed to: P. D. (email: dpx1cn@gmail.com), H. W. (email: watarih@med.hokudai.ac.jp) and J. Y. (email: jyue@uthsc.edu)



Supplementary Figure 1: Immunoblot of p53 protein expression in Hela cells after knockdown of p53.



Supplementary Figure 2: miR-34a functions as a tumor suppressor in CC cells. Invasion (left) and proliferation (right) of Hela (a) and Siha (b) cells transfected with control or miR-34a mimic. qPCR analysis of the indicated mRNAs in Hela (c) and Siha (d) cells transfected with control or miR-34a mimic. The data are shown as the mean \pm SEM; n = 3; **P< 0.01.