

**Table SI.** Number of individuals sampled per group used in this study. Complete groups for males (those used for analyses of intragroup relatedness) are those in which all adult males were sampled and complete groups for females are those in which all adult females were sampled.

Group	n M sampled	n F sampled	n immatures sampled	All M sampled?	All F sampled?
<b><i>A. pigra</i> Central</b>					
1	2	3	3	Y	Y
2	2	1	1	Y	N
3	2	1	1	Y	N
4	1	3	1	N	Y
5	3	3	1	Y	Y
10A	1	2	0	N	Y
20A	2	1	1	Y	N
10	2	3	2	Y	Y
11	2	2	1	Y	Y
12	1	2	1	N	Y
13	2	1	1	Y	N
<b><i>A. pigra</i> Outlier</b>					
C	2	2	1	N	Y
W	1	2	1	N	Y
<b>Total</b>	23	26	15	8 complete groups	9 complete groups
<b><i>A. palliata</i> West</b>					
A	1	4	0	N	Y
J	1	1	0	N	N
K	1	0	1	N	N
R	1	2	2	N	Y
Y	1	3	2	N	Y
78	3	1	1	Y	N
79	4	1	1	N	N
<b><i>A. palliata</i> East</b>					
69	1	2	2	N	N
74	3	10	9	Y	N
77	2	1	0	Y	N
80	2	2	2	Y	N
81	1	2	3	N	N
B	1	3	1	N	Y
14	2	5	0	Y	N
22	2	2	0	N	N
24	1	1	0	N	N

25	2	4	0	Y	Y
26	2	1	1	Y	N
42	3	8	3	N	N
43	0	0	1	N	N
45	4	0	3	N	N
49	1	3	3	N	N
52	2	1	1	N	N
53	1	2	3	N	Y
<b>Total</b>	42	59	39	7 complete groups	6 complete groups

---

M=adult males, F=adult females