

**Table S3. Distribution of the urinary BPA, BPF and ClxBPA levels between the two countries.**

	<b>Cyprus</b>	<b>Romania</b>	<b>p-value</b>
<b>N</b>	122	90	
<b>BPA (ng/L)</b>	1508 [789, 2806]	3778 [2134, 9050]	<0.001
<b>BPF (ng/L)</b>	485 [365, 680]	416 [219, 822]	0.091
<b>ClxBPA (ng/L)</b>	152 [141.00, 168.00]	177 [158, 201]	<0.001
<b>Creatinine-adjusted concentrations</b>			
<b>BPA (ng/g)</b>	2101 [1184, 4220]	4670 [2231, 10050]	<0.001
<b>BPF (ng/g)</b>	646 [398, 1327]	500 [271, 1138]	0.014
<b>ClxBPA (ng/g)</b>	199.19 [120, 376]	201 [125, 394]	0.961

Median[IQR] values are presented along with the *p*-value of the Wilcoxon test.

Abbreviations: BPA: bisphenol A, BPF: bisphenol F; ClxBPA: sum of 3-chlorobisphenol A (ClBPA), 3,5-dichlorobisphenol A (3,5-Cl<sub>2</sub>BPA) and 3,3'-dichlorobisphenol A (3,3'-Cl<sub>2</sub>BPA)

.