

Fibroblast miR-210 overexpression is independently associated with clinical failure in Prostate Cancer – a multicenter (*in situ* hybridization) study

Sigve Andersen, PhD, M.D.^{1,2}

Elin Richardsen, PhD, M.D.^{3,4}

Line Moi PhD, M.D.^{3,4}

Tom Donnem, PhD, M.D.^{1,2}

Yngve Nordby, M.D.^{1,5}

Nora Ness,³

Marte Eilertsen Holman, PhD, M.D.^{1,2}

Roy M. Bremnes, PhD, M.D.^{1,2}

Lill-Tove Busund, PhD, M.D.^{3,4}

¹Translational Cancer Research Group, Dept Clinical Medicine, UiT, The Arctic University of Norway

²Dept Oncology, University Hospital of North Norway, Tromso

³Translational Cancer Research Group, Dept of Medical Biology, UiT, The Arctic University of Norway

⁴Dept Pathology, University Hospital of North Norway, Tromso

⁵Dept of Urology, University Hospital of North Norway, Tromso

Corresponding author:

Sigve Andersen, PhD, MD

Department of Oncology



University Hospital of North Norway

9038 Tromso, Norway

Telephone: +47 77 62 60 00 Fax: +47 77 62 67 79

E-mail: sigve.andersen@uit.no

Staining controls of miR-210



400x microscope pictures of a) negative control of miR-210 staining (scramble) and b) positive control of miR-210 staining (U6).