

S2 Table. Bacterial meningitis epidemiology in infants aged <1 month and persons aged ≥65 years, Burkina Faso, 2011–2013

	<1 month N (%)	≥65 years N (%)
Suspected meningitis cases	15	149
Reported deaths	1 (7)	36 (24)
Probable bacterial meningitis cases	7 (47)	78 (52)
Laboratory-confirmed meningitis cases ^a	4 (27)	34 (23)
<i>H. influenzae</i>	0 (0)	0 (0)
<i>N. meningitidis</i>	1 (25)	5 (15)
Reported deaths	0 (0)	0 (0)
Meningococcal serogroup		
W	1 (100)	2 (67)
X	0 (0)	1 (33)
Missing serogroup	0	2
<i>S. pneumoniae</i>	3 (75)	29 (85)
Reported deaths	1 (33)	14 (48)
Pneumococcal serotype		
1	1 (100)	6 (30)
4	0 (0)	1 (5)
7F/7A	0 (0)	2 (10)
12F/12A/12B/44/46	0 (0)	4 (20)
19A	0 (0)	1 (5)
23F	0 (0)	1 (5)
Non-typeable	0 (0)	5 (25)
Missing serotype	2	9

^a A confirmed case of meningitis is a suspected or probable case with *S. pneumoniae*, *N. meningitidis*, or *H. influenzae* isolated from CSF by culture or detected in cerebrospinal fluid by real-time polymerase chain reaction or latex agglutination.