

1 **Employing *Escherichia coli*-derived outer membrane vesicles as an**
2 **antigen delivery platform elicits protective immunity against**
3 ***Acinetobacter baumannii* infection**

4

5 **Weiwei Huang^{1,2,3}, Shijie Wang^{1,2,3}, Yufeng Yao^{1,2,3}, Ye Xia^{1,2,3}, Xu Yang^{1,2,3}, Kui Li^{1,2,3}, Pengyan**
6 **Sun^{1,2,3}, Cunbao Liu^{1,2,3}, Wenjia Sun^{1,2,3}, Hongmei Bai^{1,2,3}, Xiaojie Chu^{1,2,3}, Yang Li^{1,2,3}, Yanbing**
7 **Ma^{1,2,3*}**

8 ¹ Laboratory of Molecular Immunology, Institute of Medical Biology, Chinese Academy of Medical
9 Sciences & Peking Union Medical College; Kunming, China 650118, ²Yunnan Key Laboratory of Vaccine
10 Research & Development on Severe Infectious Diseases; Kunming, China 650118, ³Yunnan Engineering
11 Research Center of Vaccine Research and Development on Severe Infectious Diseases, Kunming,
12 China 650118.

13

14 ***Corresponding:**

15 Prof. Yanbing Ma

16 Laboratory of Molecular Immunology,

17 Institute of Medical Biology,

18 Chinese Academy of Medical Sciences & Peking Union Medical College,

19 Kunming, China 650118

20 Phone: +86 871 68339287

21 Fax: +86 871 68334483

22 Email: yanbingma1969@126.com

23 **Supplementary figure legends**

24 **Supplementary figure 1. The DNA sequencing graph of ClyA gene**

25 DNA sequencing was used to analyze the ClyA gene from a clinical pathogenic *E.coli* W-15 and
26 non-pathogenic *E.coli* DH5 α , the result shows the ClyA gene from the DH5 α strain lacked a C base at
27 position 217 and an A base at position 493 (Black arrow).

28

29 **Supplementary Figure 1**

30

