

SUPPLEMENTARY INFORMATION

Activation of FXR protects against renal fibrosis via suppressing Smad3 expression

Kai Zhao^{a,b,†}, Jialin He^{a,†}, Yan Zhang^a, Zhizhen Xu^a, Haojun Xiong^a, Rujun Gong^c, Song Li^d, Shan Chen^{a,*}, and Fengtian He^{a,*}


^aDepartment of Biochemistry and Molecular Biology, College of Basic Medical Sciences, Third Military Medical University, Chongqing 400038, China. ^bDepartment of Pathogenic Biology, College of Basic Medical Sciences, Third Military Medical University, Chongqing 400038, China. ^cDivision of Kidney Disease and Hypertension, Department of Medicine, Rhode Island Hospital, Brown University School of Medicine, Providence, RI 02903, USA. ^dDepartment of Pharmaceutical Sciences, School of Pharmacy, University of Pittsburgh, Pittsburgh, PA 15261, USA. Correspondence and requests for materials should be addressed to F.H. (email: hefengtian06@aliyun.com) or S.C. (email: cs723cn@hotmail.com)

Shan Chen: Tel: +862368752258, E-mail: cs723cn@hotmail.com

Fengtian He: Tel./fax: +862368752255, E-mail: hefengtian06@aliyun.com

Department of Biochemistry and Molecular Biology, College of Basic Medical Sciences, Third Military Medical University, 30 Gaotanyan, Shapingba, Chongqing 400038, China.

Supplementary Figure S1. The correlation of FXR and Smad2 in human fibrotic kidneys. The expression of FXR and Smad2 in 15 human fibrotic kidneys and 15 normal control kidney tissues were examined by qRT-PCR, taking β -actin as a control. The correlation between the levels of FXR and Smad2 in human fibrotic kidneys were analyzed with Pearson's test (FXR/Smad4 $r = -0.4332$, $P = 0.1067$).



Supplementary Table S1. Subject demographics

	Control group (n=15)	Fibrosis group (n=15)
Female	3	6
Male	12	9
Average age (year)	59.6	50.3
Age(min)	35	24
Age(max)	72	80

Supplementary Table S2. The primer sets for qRT-PCR

Gene	Product size (bp)	Forward primers	Reverse primers
hFN	180	5'-tgtaccatcgcaaaccg-3'	5'-ggaagtcccagcagcat-3'
hFXR	165	5'-tacatgcgaagaaagtgtcaaga-3'	5'-actgtcttcattcacggctgtat-3'
hSHP	170	5'-cccaagatgtgtgacctt-3'	5'-ccagaaggactccagacagc-3'
hSmad2	182	5'-cggcggagaagca-3'	5'-ttaccaaaggcagcaa-3'
hSmad3	120	5'-gagtgcctcagtgacagc-3'	5'-ggtagacagcctcaaagc-3'
hSmad4	137	5'-gaacattggatgggagg-3'	5'-taagtcaaacgcatactga-3'
h β -actin	157	5'-gtgaaggtgacagcagtcggtt-3'	5'-gaagtgggtggtttagga-3'
mFN	151	5'-gaggcaggetcagcaaat -3'	5'-tccatagcaggtacaacc -3'
mFXR	148	5'-tgtgagggctgcaaaggtt-3'	5'-acatccccatctttgcac-3'
mSmad2	107	5'-gaaagccatcaccactca-3'	5'-tgtcccactgtatctaccg-3'
mSmad3	193	5'-gagaagtggtgcgagaag-3'	5'-acaggcggcagtagataa-3'
mSmad4	187	5'-agaacattggatggacgact-3'	5'-agacaacccgctcatagt-3'
m β -actin	165	5'-tgttaccaactggacgaca-3'	5'-gggtgttgaaggctcaaa-3'

Abbreviation: h, human; m, mouse.