

**Supplemental Table 1:** Cut-off values according to the manufacturers' recommendations

	IgG (IU/mL)			IgM		
	Negative	Equivocal	Positive	Negative	Equivocal	Positive
Advia						
Centaur <sup>®2</sup>	<6.4	6.4-10	≥10	<0.9	0.9-1	≥1
Architect <sup>®2</sup>	<1.5	1.6-3	≥3	<0.5	0.5-0.6	≥0.6
AxSYM <sup>®1</sup>	<2	2-3	≥3	<0.5	0.55-0.6	≥0.6
Elecsys <sup>®2</sup>	<1	1-30	≥30	<0.8	0.8-1	≥1
Enzygnost <sup>®</sup>	<4	4-10	≥10	*	*	*
Liaison <sup>®2</sup>	<7.2	7.2-8.8	≥8.8	<6	6-8	≥8
Platelia <sup>®2</sup>	<6	6-9	≥9	<0.8	0.8-1	≥1
VIDAS <sup>®1</sup>	<4	4-8	≥8	<0.55	0.55-0.65	≥0.65
VIDIA <sup>®1</sup>	<2	3-5	≥5	<0.68	0.68-1	≥1

<sup>1</sup>: Assessment 1

<sup>2</sup>: Assessment 2

\* In each series, the cut-off value was calculated by adding 0.200 to the mean of the absorbance values of Toxoplasmosis/IgM negative control serum. After application of a correction factor (calculated from the Toxoplasmosis/IgM positive control serum) on the absorbance values, the resulting value "A" was interpreted as follows: negative if  $A < \text{cut-off}$ , positive if  $A > \text{cut-off} + 0.050$ , equivocal if  $\text{cut-off} \leq A \leq \text{cut-off} + 0.050$