Rebholz et al, AJKD, "DASH (Dietary Approaches to Stop Hypertension) Diet and Risk of Subsequent Kidney Disease"

Table S5. Risk of Kidney Disease by Tertile of the Alternative DASH Diet Score Based on Nutrients

Tertile of the Alternative DASH Diet Score (Nutrients)				
	Tertile 1 (0-2.5):	Tertile 2 (3-4):	Tertile 3 (4.5-9):	P-value
	Low Score	Moderate Score	High Score	for trend
Model 1	1.06 (0.97, 1.15)	0.96 (0.89, 1.05)	1 [Reference]	0.16
Model 2	1.04 (0.95, 1.14)	0.93 (0.86, 1.01)	1 [Reference]	0.26
Model 3	1.11 (1.02, 1.22)	0.94 (0.87, 1.03)	1 [Reference]	0.007

Model 1: Adjusted for age, sex, race-center, education level, smoking status, physical activity, total caloric intake

Model 2: Model 1 + baseline eGFR (linear spline terms with one knot at 90 mL/min/1.73 m<sup>2</sup>)

Model 3: Model 2 + overweight/obese status, diabetes, hypertension, systolic blood pressure, use of angiotensin converting enzyme inhibitors or angiotensin receptor blockers