


Developing a core outcome set for studies in weight loss surgery

Round two questionnaire



Completing the survey

Thank you for returning the first round of the survey and for continuing to take part in this study. Everyone's opinions from the first round were pooled. This second round of the survey includes the same list of questions as the first round but also includes the results of the first round. The ratings from round one are provided for each question. These include the overall patient group rating (pink column), the overall healthcare professionals (HCPs) group rating (yellow column), and your own rating (blue column). We are now asking you to re-rate each question taking into account other people's views.

Please answer all of the questions yourself. This should take you a maximum of 30 minutes. **Don't spend too long on the questions, we are looking for your immediate feelings. The information that you provide will remain anonymous.** When you have completed the questionnaire, please return it in the pre-paid envelope provided. Thank you again so much for all your time and help with this survey, which will help us with future research.

SECTION 1 Medical effectiveness of weight loss surgery

The following section lists different ways to measure the medical effectiveness of weight loss surgery. Please rate how important you think it is that the following information is collected to measure the impact/effectiveness of the surgery and circle the number that represents your opinion. The words in brackets are the medical terminology.

HCPs = Healthcare professionals.

		Average previous ratings											
		Patients	HCPs	Your own	Not important			(Please circle)			Extremely important		
For example: Less pain when walking		4	5	6	1	2	3	4	5	6	7	8	9
		Patients	HCPs	Your own	Not important			(Please circle)			Extremely important		
Body measurements (anthropometry)													
1	A measurement of weight				1	2	3	4	5	6	7	8	9
2	Body dimensions such as waist and hip measurements				1	2	3	4	5	6	7	8	9
3	Body mass index				1	2	3	4	5	6	7	8	9
Obesity related disease													
4	Improvement in abnormal or irregular heartbeat (arrhythmia)				1	2	3	4	5	6	7	8	9
5	Reduction/lowering of blood pressure to a healthy level, or a reduction in blood pressure medication (hypertension)				1	2	3	4	5	6	7	8	9
6	Reduction in the chance of having heart problems in the future (adjusted cardiovascular risk)				1	2	3	4	5	6	7	8	9
7	Improvement in diabetes, diabetes no longer being present, or a reduction in diabetic medication (measure of diabetes e.g. HbA1c)				1	2	3	4	5	6	7	8	9

SECTION 2 Short and long term medical complications

This section lists events that may occur during, or after, weight loss surgery. Please note, these are only possibilities and do not occur in everyone. Some of these events are extremely rare. The words in brackets are the medical terminology. Please rate how important you think it is that the following events are measured in studies of weight loss surgery and circle the number that best represents your opinion.

		Average previous ratings											
Complications that might occur during the operation		Patients	HCPs	Your own	Not important			(Please circle)			Extremely important		
11	Bleeding problems, such as a collection of blood or loss of blood during the operation (peri-operative haemorrhage)				1	2	3	4	5	6	7	8	9
12	The risk of accidentally damaging other organs during the operation (organ injury)				1	2	3	4	5	6	7	8	9
13	The likelihood of needing to use a large cut/incision during surgery, rather than planned keyhole surgery (conversion to open surgery)				1	2	3	4	5	6	7	8	9
Infection													
14	Infection inside the body where the operation was carried out (deep abscess)				1	2	3	4	5	6	7	8	9
15	Problems with the wound such as infection, oozing, or failure to heal properly (wound infection or dehiscence)				1	2	3	4	5	6	7	8	9
16	Whole body infection which requires prolonged admission to hospital (septicaemia)				1	2	3	4	5	6	7	8	9

Average previous ratings

		Average previous ratings			Not important (Please circle) Extremely important								
		Patients	HCPs	Your own									
Haemorrhage													
17	Bleeding problems in stomach or bowel which may cause blood to be seen in bowel movements (gastrointestinal bleeding)				1	2	3	4	5	6	7	8	9
18	Bleeding inside the abdomen where the operation was carried out (intra-abdominal bleeding)				1	2	3	4	5	6	7	8	9
19	Bleeding from the wound (wound bleeding / port site haematoma)				1	2	3	4	5	6	7	8	9
Surgical joins between internal organs (anastomosis)													
20	Leaking of stomach contents through a hole in the stomach (gastric fistula)				1	2	3	4	5	6	7	8	9
21	Leaking of bowel contents into the abdomen through a hole where the bowel is joined or stapled (anastomotic leak)				1	2	3	4	5	6	7	8	9
22	Abnormal narrowing of the bowel caused by scar tissue or stapling, which might cause a blockage (stenosis)				1	2	3	4	5	6	7	8	9
23	Bleeding from the internal bowel staples (staple line bleed)				1	2	3	4	5	6	7	8	9
24	Ulcers developing at the new join between the two pieces of bowel (anastomotic ulceration)				1	2	3	4	5	6	7	8	9
Band related complications													
25	Infection of the gastric band (band infection)				1	2	3	4	5	6	7	8	9
26	Problems with the port, such as flipping or needing the port to be relocated (port erosion or revisions)				1	2	3	4	5	6	7	8	9

A

Average previous ratings

		Patients	HCPs	Your own	Not important (Please circle) Extremely important								
27	Infection of the port which is used to change the band size (port infection)				1	2	3	4	5	6	7	8	9
28	The gastric band eroding/growing into the stomach (band moves from outside to the inside of stomach) leading to the need for further surgery (band erosion)				1	2	3	4	5	6	7	8	9
29	The band slipping out of place and needing more surgery to correct it (band slippage)				1	2	3	4	5	6	7	8	9
30	Intolerance of the band				1	2	3	4	5	6	7	8	9
31	Pain in the stomach caused by overeating/too much food stretching the stomach above the band (pouch dilation)				1	2	3	4	5	6	7	8	9
32	Damage to the band, which might require band replacement (iatrogenic injury [device])				1	2	3	4	5	6	7	8	9
33	Failure of the port or tube to work properly (port malfunction)				1	2	3	4	5	6	7	8	9
34	The need for further surgery to make changes to the band (band revisions)				1	2	3	4	5	6	7	8	9
Obstruction / hernia													
35	Twisting or abnormal movement of the bowel or intestines, which can cause blockages, pain or nausea and may need additional surgery (internal hernia)				1	2	3	4	5	6	7	8	9
36	A lump or bulge in the skin through the scar where the surgery was performed (external hernia, incisional)				1	2	3	4	5	6	7	8	9

Average previous ratings

		Patients	HCPs	Your own	Not important (Please circle) Extremely important								
37	Build up of scar tissue in the bowel causing obstruction or abdominal pain (adhesional obstruction)				1	2	3	4	5	6	7	8	9
38	Vomiting or excessive bloating after the operation whilst still in hospital (ileus)				1	2	3	4	5	6	7	8	9

General complications of surgery

39	One or more areas of the lungs collapsing or not inflating properly (atelectasis)				1	2	3	4	5	6	7	8	9
40	Needing a machine to help with breathing (ventilation)				1	2	3	4	5	6	7	8	9
41	Chest infection (lower respiratory tract infection)				1	2	3	4	5	6	7	8	9
42	Heart's blood supply is blocked, or interrupted, by a build-up of fatty substances in the heart's arteries (ischaemic/coronary heart disease)				1	2	3	4	5	6	7	8	9
43	Irregular or abnormal heart beat (arrhythmia)				1	2	3	4	5	6	7	8	9
44	Blood clot in the leg or lung (venous thromboembolism)				1	2	3	4	5	6	7	8	9
45	Stroke (cerebrovascular accident)				1	2	3	4	5	6	7	8	9
46	Kidney failure (renal failure)				1	2	3	4	5	6	7	8	9
47	Bladder infection in 'wee/pee' (Urinary tract infection)				1	2	3	4	5	6	7	8	9
48	Unintentional passing of urine (incontinence)				1	2	3	4	5	6	7	8	9

Mortality

49	Risk of death during the operation (peri-operative mortality)				1	2	3	4	5	6	7	8	9
----	---	--	--	--	---	---	---	---	---	---	---	---	---

SECTION 3 Physical signs, symptoms and other measures

The following section lists some physical signs and symptoms that some people may experience before or after weight loss surgery. It also lists some measurements that may be taken. Some of these signs and symptoms may get better after surgery. Some may get worse, or remain the same. Please note, these are only possibilities and do not occur in everyone. The words in brackets are the medical terminology.

Please rate how important you think it is that the following symptoms are measured in research studies of weight loss surgery, and circle the number that best represents your opinion.

		Average previous ratings											
Signs		Patients	HCPs	Your own	Not important			(Please circle)			Extremely important		
53	Hair loss				1	2	3	4	5	6	7	8	9
54	Problems hearing				1	2	3	4	5	6	7	8	9
55	Problems with gums or teeth				1	2	3	4	5	6	7	8	9
56	Problems with vision				1	2	3	4	5	6	7	8	9
57	Swelling or retaining water (oedema)				1	2	3	4	5	6	7	8	9
58	Skin problems or irritations (such as rashes, sores or loose skin, ulcers)				1	2	3	4	5	6	7	8	9
59	Loss of sensation in hands and feet (peripheral neuropathy/paraesthesia)				1	2	3	4	5	6	7	8	9
Symptoms													
60	Feeling out of breath (breathlessness)				1	2	3	4	5	6	7	8	9
61	Belching, bloating, or gas (flatulence)				1	2	3	4	5	6	7	8	9
62	Constipation or difficulty passing stool				1	2	3	4	5	6	7	8	9
63	Diarrhoea or loose bowel motion				1	2	3	4	5	6	7	8	9

Average previous ratings

		Patients	HCPs	Your own	Not important (Please circle) Extremely important								
64	Pain or discomfort in the body				1	2	3	4	5	6	7	8	9
65	Feeling hot or sweaty				1	2	3	4	5	6	7	8	9
66	Feeling light-headed or dizzy				1	2	3	4	5	6	7	8	9
67	Feeling sick or vomiting (nausea)				1	2	3	4	5	6	7	8	9
68	Numbness or tingling in the body				1	2	3	4	5	6	7	8	9
69	Heartburn or acid indigestion (reflux)				1	2	3	4	5	6	7	8	9
70	Problems swallowing or bringing food back up (dysphagia/regurgitation)				1	2	3	4	5	6	7	8	9
71	Problems controlling the bladder (urinary incontinence)				1	2	3	4	5	6	7	8	9
72	Appearing physically tired or lacking in energy (physical fatigue)				1	2	3	4	5	6	7	8	9
73	Pain or discomfort in stomach area				1	2	3	4	5	6	7	8	9
Other Measures													
74	A measurement of <u>vitamin</u> levels				1	2	3	4	5	6	7	8	9
75	How many calories consumed (energy intake)				1	2	3	4	5	6	7	8	9
76	A measurement of <u>mineral</u> levels				1	2	3	4	5	6	7	8	9
77	The length of time spent in hospital after admission for surgery (length of hospital stay)				1	2	3	4	5	6	7	8	9
78	How long the operation takes (operative time)				1	2	3	4	5	6	7	8	9
79	Unexpected return to hospital for unplanned procedures or urgent review (re-admission rates)				1	2	3	4	5	6	7	8	9

SECTION 4 Impact of surgery on quality of life and wellbeing

The following section lists some other areas of life that can be affected by having weight loss surgery. Please rate how important you think it is, that the following information is measured in research studies of weight loss surgery, and circle the number that best represents your opinion.

		Average previous ratings											
Activities of daily living and work/employment		Patients	HCPs	Your own	Not important (Please circle) Extremely important								
		1 2 3 4 5 6 7 8 9											
80	Being able to carry out usual activities (not related to paid employment) such as personal hygiene, housework, managing finances				1	2	3	4	5	6	7	8	9
81	Being able to shop for clothes that fit				1	2	3	4	5	6	7	8	9
82	Being able to fit into spaces in public places (e.g. fit into seats or through aisles)				1	2	3	4	5	6	7	8	9
83	Mobility (e.g. being able to walk, climb stairs, bend, cross legs, get up from chairs)				1	2	3	4	5	6	7	8	9
84	Fitness (strength and endurance)				1	2	3	4	5	6	7	8	9
85	Being able to participate in, and enjoy physical activities				1	2	3	4	5	6	7	8	9
86	Being able to accomplish work tasks, or to take up work/paid employment				1	2	3	4	5	6	7	8	9
87	Satisfaction and recognition at work (if in paid employment)				1	2	3	4	5	6	7	8	9
88	Relationships with work colleagues (if in paid employment)				1	2	3	4	5	6	7	8	9
Body image													
89	Feeling satisfied and confident with one's body				1	2	3	4	5	6	7	8	9

Average previous ratings

		Patients	HCPs	Your own	Not important (Please circle) Extremely important								
90	Feeling in control of weight and appearance				1	2	3	4	5	6	7	8	9
91	Feeling like the mind and body are in tune with each other				1	2	3	4	5	6	7	8	9
92	Excess skin or skin folds following weight loss				1	2	3	4	5	6	7	8	9
Eating behaviour													
93	Having a healthy/balanced eating pattern				1	2	3	4	5	6	7	8	9
94	Being able to recognise hunger feelings				1	2	3	4	5	6	7	8	9
95	Being able to stop eating when feeling full				1	2	3	4	5	6	7	8	9
96	Time spent thinking about food				1	2	3	4	5	6	7	8	9
97	Eating for emotional reasons				1	2	3	4	5	6	7	8	9
98	Feeling guilty or upset after eating				1	2	3	4	5	6	7	8	9
99	Eating in secret				1	2	3	4	5	6	7	8	9
100	Eating differently in social situations than normal				1	2	3	4	5	6	7	8	9
Psychological and emotional wellbeing													
101	How individuals feel others perceive them				1	2	3	4	5	6	7	8	9
102	Self-esteem and self-confidence (how someone perceives themselves)				1	2	3	4	5	6	7	8	9
103	Mood swings				1	2	3	4	5	6	7	8	9
104	Stress levels				1	2	3	4	5	6	7	8	9

Average previous ratings

		Patients	HCPs	Your own	Not important (Please circle) Extremely important								
105	Coping (how someone feels with stress or difficulties)				1	2	3	4	5	6	7	8	9
106	Feeling in control of emotional or psychological wellbeing				1	2	3	4	5	6	7	8	9
Mental health													
107	Depression				1	2	3	4	5	6	7	8	9
108	Anxiety				1	2	3	4	5	6	7	8	9
109	Hostility (level of anger, and ability to manage this)				1	2	3	4	5	6	7	8	9
110	Fears or phobias				1	2	3	4	5	6	7	8	9
111	Suicidal thoughts				1	2	3	4	5	6	7	8	9
112	Other addictive behaviours (e.g. alcohol, drugs, gambling, shopping)				1	2	3	4	5	6	7	8	9
Sex life													
113	Interest in and enjoyment of sex				1	2	3	4	5	6	7	8	9
114	Ability to physically participate in sex				1	2	3	4	5	6	7	8	9
115	Sexual confidence				1	2	3	4	5	6	7	8	9
116	Sexual attention from others				1	2	3	4	5	6	7	8	9
Sleep													
117	Ability to fall asleep at night				1	2	3	4	5	6	7	8	9
118	Overall quality of sleep				1	2	3	4	5	6	7	8	9
119	Sleepiness during the day				1	2	3	4	5	6	7	8	9
120	Snoring which affects others				1	2	3	4	5	6	7	8	9

Average previous ratings

		Average previous ratings			Not important (Please circle) Extremely important									
		Patients	HCPs	Your own	1	2	3	4	5	6	7	8	9	
Social														
121	Relationship with partner/spouse				1	2	3	4	5	6	7	8	9	
122	Relationship with, and/or ability to care for, children				1	2	3	4	5	6	7	8	9	
123	Relationship with other family members				1	2	3	4	5	6	7	8	9	
124	Relationship with friends				1	2	3	4	5	6	7	8	9	
125	Treatment from people in wider society				1	2	3	4	5	6	7	8	9	
126	Having confidence to participate in social activities				1	2	3	4	5	6	7	8	9	
127	Feeling in control of social life				1	2	3	4	5	6	7	8	9	
Overall health, wellbeing and life														
128	Normality (feeling able to live a 'normal' life)				1	2	3	4	5	6	7	8	9	
129	Feeling in control of health and wellbeing				1	2	3	4	5	6	7	8	9	
130	Having a positive outlook on life and expectations for the future				1	2	3	4	5	6	7	8	9	

Thank you for your contribution to this round of the survey. Please now return the questionnaire in the pre-paid envelope provided.

Contact details
 Freepost RTKR-UYTS-AKER
 University of Bristol
 School of Social & Community Medicine
 Canynge Hall
 39 Whatley Road
 Bristol
 BS8 2PS

 karen.coulman@bristol.ac.uk