

Appendix 2 -Prevalence of risk behaviors (%) among Brazilian adolescents. National Survey of Health among 9th Grade Students (PeNSE, 2012)

Behaviors (%)	Gender	Operationalization						
		no	1-2 days	3-5 days	6-9 days	10-19 days	20-29 days	everyday
Substance use								
Smoking - last 30 days	Male	94.9	2.6	0.6	0.5	0.4	0.2	0.7
	Female	95.0	2.8	0.7	0.4	0.4	0.2	0.5
	Total	94.9	2.7	0.7	0.5	0.4	0.2	0.6
Drinking alcohol - last 30 days	Male	74.8	15.1	4.4	2.4	1.8	0.9	0.7
	Female	73.1	16.7	4.6	2.4	1.8	1.0	0.4
	Total	73.9	16.0	4.5	2.4	1.8	1.0	0.6
Amount of alcohol (Glasses)								
	Male	72.8	< 1 glass	1 glass	2 glasses	3 glasses	4 glasses	>=5 glasses
	Female	69.6	7.2	7.1	3.7	2.6	1.7	4.9
	Total	71.2	8.4	7.6	3.9	2.6	1.8	4.6
Drug use - last 30 days								
	Male	97.2	1-2 days	3-5 days	6-9 days	>=10 days		
	Female	98.0	1.3	0.5	0.3	0.7		
	Total	97.6	1.2	0.3	0.2	0.3		
Sexual behavior								
Number of sexual partners	Male	59.9	1	2	3	4	5	6 or more
	Female	81.7	13.9	8.2	5.0	2.9	1.9	8.2
	Total	71.3	10.6	3.0	1.7	0.9	0.7	1.4
Unsafe sex (without condom)	Male	90.7	12.2	5.5	3.2	1.9	1.3	4.7
	Female	94.8	no	yes				

		Total	7.2	2-3 times	4-5 times	6-7 times	8-9 times	10-11 times	12 or more times
Aggression		92.8	once						
		no							
Physical fight	Male	72.7	13.5	7.2	2.6	1.1	0.7	0.3	2.0
	Female	85.4	8.8	3.5	1.0	0.3	0.3	0.1	0.6
	Total	79.3	11.0	5.3	1.8	0.7	0.5	0.2	1.3
		no	yes						
Bullying	Male	73.9	26.2						
	Female	84.0	16.0						
	Total	79.2	20.8						
Fight using gun	Male	91.3	8.8						
	Female	95.7	4.3						
	Total	93.6	6.4						
Fight using other weapons	Male	89.9	10.1						
	Female	95.2	4.8						
	Total	92.7	7.3						
