Injections of intravenous contrast for computerized tomography scans precipitate migraines in hereditary hemorrhagic telangiectasia subjects at risk of paradoxical emboli: implications for right-to-left shunt risks.

Patel T, Elphick A, Jackson JE, Shovlin CL

SUPPLMENTARY DATA

Extracts from the questionnaire designed in SurveyMonkey¹ at https://www1.imperial.ac.uk/nhli/cardio/vascular/pulmonary_vascular/hht_pavms/hhtsurvey/

¹SurveyMonkey.com, LLC, Palo Alto, California, USA (www.surveymonkey.com)

Imperial College London

Supporting HHT families
HHTIC London
through clinical care and research

12/EM/0073: HHT and other medical conditions (2013)

We invite you to help us with this research, by filling out the enclosed short questionnaire about general health issues. There are 5 sections, and it should take you about 20-30 minutes to fill in. This will allow us to see if people with HHT are more or less likely to have certain common medical conditions, than people without HHT. All questionnaires will be treated with the strictest of confidence. We will not disclose your personal results to your family doctor, or any third party.

It is up to you to decide whether to fill in the questionnaire. If you do fill it in and press the "DONE" button on the last page, we will accept your consent to use your answers in our research. We aim to publish our conclusions in a scientific journal. No study participant will be identified in any publication. Thank you very much for considering this invitation.

Dr Claire Shovlin, Lead Clinician, HHTIC London

Note that to reduce bias, participants were provided with very general information in their information sheet, and the online survey introduction.

lm	perial	College
	ndon	100

Supporting HHT families HHTIC London through clinical care and research

12/EM/0073: HHT and other medical conditions (2013)

47. How often do you have migraines?
I have never had a migraine
Less than 5 times in my life
About once a year
At least once a month
At least once a week
Most days
Every day
48. My migraines started when I was
younger than 5
<u> </u>
<u> </u>
in late teens
in my 20s
in my 30s
in my 40s
in my 50s
in my 60s
in my 70s
in my 80s
in my 90s
49and as I have got older, my migraines (tick all that apply)
have stayed about the same
are less frequent
are more frequent
seem less severe
seem more severe

lm	perial	College
In	ndon	

Supporting HHT families
HHTIC London
through clinical care and research

12/EM/0073: HHT and other medical conditions (2013)

5	0. What does your doctor say about your migraines?
	I have never needed to discuss them with a doctor
	My doctor says the headaches are due to something else, but I think they are migraines
	My doctor thinks my headaches are migraines, and has told me to take simple painkillers like paracetamol
	My doctor thinks my headaches are migraines, and has given me specific treatment for migraines
ŧ	51. Which of these applies to your migraines. <i>Tick all that apply</i> . The headache is
	mild in intensity
	mild to moderate in intensity
	moderate to severe in intensity
	distracting but not debilitating and I can do most of my usual activities
	a steady ache
	intense, pounding, throbbing or debilitating
	on one side of the head
	on both sides of the head
* 5	2. Does anything else happen before, during or after the headache?
	Nothing really, it's just a headache
	Yes, something else sometimes or usually happens
	,

Imperial College London supporting HHT families HHTIC London through clinical care and research 12/EM/0073: HHT and other m	nedical condition	ns (2013)			
53. You have told us that wh more, so please us how ofte		-	-	• •	
, .	Never	With some	Often	With most	Always
I feel sick or vomit	\circ	0	\circ	0	\circ
I am sensitive to bright lights	\circ	\circ	\circ	\circ	\circ
I am sensitive to loud noises		\circ	\bigcirc	\circ	\circ
I pass lots of urine	\bigcirc	\circ	\bigcirc	\bigcirc	\circ
I get tummy pains	\circ	\circ	\circ	\circ	\circ
I get diarrhoea	\bigcirc	\circ	\bigcirc	\bigcirc	\circ
I have a nosebleed	\circ	\circ	\circ	\circ	\circ
Remember, this is something that					
54. When you have had a m things), or movement of an		ou ever had a problem v	with your vision, h	earing, speech, sens	ation (how you feel
Yes					
○ No					
i'm not sure					

Imperial College London					
Supporting HHT families HHTIC London through clinical care and research					
12/EM/0073: HHT and other	medical condition	ons (2013)			
55. You have told us that w disturbance. We would like after your migraine headaci	to know a little r				
	Never	With some	Often	With most	Always
I see flashes of light	\bigcirc	\circ	\circ	\circ	\circ
Objects seem to move around in pieces	\circ	\circ	\circ	\circ	\circ
I get a temporary loss of vision	\circ	0	\circ	\circ	0
I get a temporary loss of sensation	\bigcirc	\bigcirc	\bigcirc	\bigcirc	\circ
I have a speech disturbance	\circ	\bigcirc	0	\circ	\circ
I have a problem with movement	\bigcirc	\bigcirc	\bigcirc	\bigcirc	\circ
I have a problem with balance	\circ	\circ	\circ	\circ	\circ
You have 3 lines to give a little mo	ore detail, if you war	nt to.			

Remember, we are asking about something that happens at the time you have a migraine. We are not asking about other times in your life.

Imperial College London supporting HHT familles HHTIC London through clinical care and research						
12/EM/0073: HHT and other 121. This is a list of simple them seemed to make a decision of the seeme	e tests in hospita	ıls. We would lik	e to know if you l	have had any of t	hese tests, and i	f so, if any of
	I have not had this investigation	Migraines stayed about the same	Migraines seemed to get better	Migraines seemed to get worse	This seemed to start a migraine	This seemed to stop a migraine
Getting weighed						
Having blood pressure measured						
Having oxygen measurements by a finger probe						
Blood tests						
Giving a urine sample						
Breathing tests - blowing out hard						
Breathing tests with a nose clip						
A standard x-ray of the chest, head, abdomen or another part of the body						
You have 3 linew if you want to	give us any more de	tails.				
* 122. Have you had a CT	scan? This is a c	quiet, quick scar	n and you lie on a	bed that goes th	rough a doughn	ut-shaped ring.
○ No						
Yes						
l'm not sure			ı	f they answered	d "Yes" they we	ere directed to:

Patel T, Elphick A, Jackson JE, Shovlin CL. Injections of intravenous contrast for computerized tomography scans precipitate migraines in hereditary hemorrhagic telangiectasia subjects at risk of paradoxical emboli: implications for right-to-left shunt risks. Headache 2016

Imperial College London Supporting HHT families HHTIC London through clinical care and research						
12/EM/0073: HHT and other	er medical conditi	ions (2013)				
123. We would like to kno	w if having a CT	scan seemed to	make a difference	e to your migrain	es.	
	I have not had one	No different really	Migraines seemed a bit better	Migraines seemed a bit worse	Seemed to bring on a migraine	Seemed to stop a migraine
CT scans with no needle/injection						
CT scans with an injection of contrast						
You have 3 lines if you want to	give us any more de	tails.				
* 124. Have you had an M	RI scan? This is	a noisy scan tha	t usually lasts 15	-20 minutes or m	ore	
○ No ○ Yes						
I'm not sure						
				If they answere	d "Yes" they w	ere directed to
125. We would like to kno	ow if having an M	RI scan seemed	to make a differe	nce to your migra	aines.	
	I have not had one	No different really	Migraines seemed a bit better	Migraines seemed a bit worse	Seemed to bring on a migraine	Seemed to stop a migraine
MRI scans with no needle/injection						
MRI scans with an injection of contrast						
You have 3 lines if you want to	give us any more de	etails.				

Imperial College London Supporting HHT families						
through clinical care and research 12/EM/0073: HHT and other	er medical condition	ons (2013)				
* 126. Have you had an ec firmly, and the pictures a			robe with jelly on	will have been m	oved over your	skin, pressing
○ No						
Yes						
I'm not sure			ı	f they answered	d "Yes" they wo	ere directed to:
127. We would like to kno	ow if having an ec	hocardiogram o	r ultrasound seer	ned to make a dif	ference to your	migraines.
	I have not had one	No different really	Migraines seemed a bit better	Migraines seemed a bit worse	Seemed to bring on a migraine	Seemed to stop a migraine
Echocardiogram with no needle/injection						
Ultrasound (usually of abdomen) with no needle/injection						
Bubble echo (with an injection)						
You have 3 lines if you want to	give us any more def	tails.				