



Preliminary Full wwPDB X-ray Structure Validation Report ⓘ

Oct 25, 2016 – 06:23 AM EDT

This is a Preliminary Full wwPDB X-ray Structure Validation Report.

This report is produced by the standalone wwPDB validation server.
**The structure in question has not been deposited to the wwPDB.
This report should not be submitted to journals.**

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.1 (RC1), CSD as537be (2016)
Xtrriage (Phenix)	:	1.9-1692
EDS	:	rb-20027939
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027939


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	281	 88% 12%
2	B	283	 89% 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	UMP	D	314	X	-	-	-
4	UMP	D	315	X	-	-	-

PRELIMINARY VALIDATION REPORT

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4632 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	281	2239	1431	399	397	12	0	0	0

- Molecule 2 is a protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	281	2242	1431	397	401	13	0	0	0

- Molecule 3 is TOMUDEX (three-letter code: D16) (formula: unknown).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	C	1	32	21	4	6	1	0	0
3	C	1	32	21	4	6	1	0	0

- Molecule 4 is 2'-DEOXYURIDINE 5'-MONOPHOSPHATE (three-letter code: UMP) (formula: unknown).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	D	1	20	9	2	8	1	0	0
4	D	1	20	9	2	8	1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	S	47	Total 47 47	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1:



- Molecule 2:



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.41Å 79.54Å 67.28Å 90.00° 116.24° 90.00°	Depositor
Resolution (Å)	19.92 – 2.40 48.08 – 2.36	Depositor EDS
% Data completeness (in resolution range)	98.6 (19.92-2.40) 95.8 (48.08-2.36)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.37 (at 2.37Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.178 , 0.232 0.179 , 0.233	Depositor DCC
R_{free} test set	1178 reflections (5.07%)	DCC
Wilson B-factor (Å ²)	55.8	Xtrriage
Anisotropy	0.109	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.020 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4632	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: D16, UMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/2299	0.58	0/3120
2	B	0.46	0/2302	0.62	0/3124
All	All	0.45	0/4601	0.60	0/6244

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2239	0	2202	23	0
2	B	2242	0	2187	23	0
3	C	64	0	40	5	0
4	D	40	0	20	3	0
5	S	47	0	0	0	0
All	All	4632	0	4449	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:84:GLY:HA2	2:B:235:GLN:HE21	1.33	0.90
2:B:235:GLN:HG2	2:B:273:ILE:HB	1.70	0.74
1:A:102:ARG:HB3	1:A:330:THR:HG22	1.72	0.71
1:A:220:HIS:HB3	1:A:236:LEU:HD11	1.76	0.68
2:B:242:ASP:HB2	4:D:315:UMP:H2"	1.75	0.66
1:A:114:TRP:HA	1:A:117:LYS:HE2	1.78	0.65
1:A:84:GLY:HA2	2:B:235:GLN:NE2	2.09	0.64
1:A:302:ARG:N	1:A:323:VAL:O	2.32	0.59
1:A:242:ASP:HB2	4:D:314:UMP:H2"	1.86	0.58
2:B:54:GLU:OE2	2:B:100:THR:OG1	2.14	0.56
2:B:220:HIS:HB2	2:B:236:LEU:HD11	1.87	0.56
3:C:414:D16:HP12	4:D:314:UMP:O4	2.06	0.56
2:B:158:VAL:HG12	2:B:159:TYR:H	1.72	0.54
1:A:121:ASP:CG	1:A:151:ARG:HH22	2.12	0.53
2:B:249:PHE:CG	3:C:415:D16:H15	2.44	0.53
1:A:104:PHE:CD2	3:C:414:D16:HB2	2.43	0.52
2:B:158:VAL:CG1	2:B:159:TYR:H	2.22	0.52
1:A:73:ASP:OD1	1:A:77:ILE:N	2.44	0.51
2:B:158:VAL:HG12	2:B:159:TYR:N	2.26	0.50
2:B:220:HIS:ND1	2:B:220:HIS:N	2.60	0.49
1:A:249:PHE:CG	3:C:414:D16:H15	2.48	0.49
1:A:208:PRO:HD2	2:B:166:PHE:CE1	2.48	0.49
2:B:55:LEU:O	2:B:59:ARG:HG3	2.14	0.48
2:B:220:HIS:CB	2:B:236:LEU:HD11	2.43	0.48
2:B:88:ARG:NH2	2:B:271:GLU:OE1	2.38	0.47
1:A:73:ASP:OD2	1:A:75:ALA:HB3	2.15	0.46
1:A:70:ASP:OD2	1:A:283:LYS:NZ	2.46	0.46
1:A:287:GLU:HB2	1:A:288:PRO:HD3	1.98	0.46
2:B:293:LEU:HA	2:B:293:LEU:HD23	1.76	0.45
1:A:247:VAL:HG13	1:A:274:HIS:HE1	1.82	0.45
2:B:292:GLN:HA	2:B:295:ARG:HD2	1.98	0.45
2:B:161:PHE:CZ	2:B:168:ALA:HB3	2.52	0.44
1:A:122:SER:OG	1:A:153:GLY:HA2	2.19	0.43
1:A:275:THR:HG21	2:B:275:THR:HG21	2.01	0.43
2:B:156:GLY:HA2	2:B:170:TYR:CE2	2.54	0.42
3:C:415:D16:H15	3:C:415:D16:HP11	1.88	0.42
2:B:158:VAL:CG1	2:B:159:TYR:N	2.83	0.42
2:B:55:LEU:HD23	2:B:58:LEU:HD12	2.02	0.41
2:B:285:HIS:NE2	2:B:333:MET:HB3	2.36	0.41
1:A:121:ASP:OD2	1:A:151:ARG:NH2	2.46	0.40
1:A:177:TYR:O	1:A:180:GLN:HB2	2.21	0.40
1:A:247:VAL:O	1:A:251:ILE:HG13	2.22	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:214:MET:SD	1:A:218:PRO:HD3	2.61	0.40
1:A:61:LEU:HA	1:A:61:LEU:HD23	1.95	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	279/281 (99%)	267 (96%)	11 (4%)	1 (0%)	39	56
2	B	277/283 (98%)	262 (95%)	14 (5%)	1 (0%)	39	56
All	All	556/564 (99%)	529 (95%)	25 (4%)	2 (0%)	39	56

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	104	PHE
1	A	158	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/234 (100%)	233 (100%)	1 (0%)	93	98
2	B	235/234 (100%)	231 (98%)	4 (2%)	68	85

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	469/468 (100%)	464 (99%)	5 (1%)	80	92

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	220	HIS
2	B	171	VAL
2	B	220	HIS
2	B	235	GLN
2	B	292	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	235	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	D16	C	414	-	33,?,?	2.49	9 (27%)	42,?,?	2.60	13 (30%)
3	D16	C	415	-	33,?,?	2.59	8 (24%)	42,?,?	2.74	12 (28%)
4	UMP	D	314	-	21,?,?	3.69	10 (47%)	30,?,?	2.80	16 (53%)
4	UMP	D	315	-	21,?,?	3.67	10 (47%)	30,?,?	2.83	11 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	D16	C	414	-	-	0/25/?/?	0/3/?/?
3	D16	C	415	-	-	0/25/?/?	0/3/?/?
4	UMP	D	314	-	1/1/?/?	0/10/?/?	0/2/?/?
4	UMP	D	315	-	1/1/?/?	0/10/?/?	0/2/?/?

All (37) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	315	UMP	C2'-C3'	-8.37	1.30	1.52
4	D	314	UMP	C2'-C3'	-8.25	1.30	1.52
4	D	314	UMP	O4'-C4'	-7.80	1.27	1.45
4	D	315	UMP	O4'-C4'	-7.46	1.27	1.45
4	D	315	UMP	C1'-N1	-4.64	1.35	1.48
4	D	314	UMP	C1'-N1	-4.62	1.35	1.48
3	C	414	D16	C14-S13	-2.13	1.70	1.73
3	C	415	D16	C9-C6	2.11	1.55	1.51
4	D	315	UMP	C6-N1	2.13	1.43	1.37
4	D	315	UMP	C4-N3	2.29	1.43	1.38
4	D	314	UMP	C6-N1	2.33	1.43	1.37
3	C	414	D16	C15-C14	2.35	1.44	1.36
4	D	314	UMP	C4-N3	2.46	1.43	1.38
3	C	415	D16	CG-CD	2.73	1.57	1.50
3	C	414	D16	CG-CD	2.77	1.57	1.50
3	C	415	D16	C5-C6	2.93	1.44	1.39
3	C	414	D16	C5-C6	3.07	1.44	1.39
3	C	415	D16	C5-C4A	3.12	1.45	1.39
3	C	414	D16	C-C11	3.27	1.53	1.47
4	D	315	UMP	C3'-C4'	3.34	1.62	1.53
3	C	414	D16	C5-C4A	3.37	1.45	1.39
4	D	314	UMP	C3'-C4'	3.42	1.62	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	415	D16	C-C11	3.78	1.54	1.47
4	D	314	UMP	C2-N3	4.34	1.46	1.38
3	C	415	D16	C8-C8A	4.53	1.47	1.39
3	C	414	D16	C8-C8A	4.62	1.47	1.39
4	D	315	UMP	C2-N3	4.70	1.46	1.38
4	D	315	UMP	C2-N1	4.89	1.46	1.38
4	D	314	UMP	C2-N1	4.92	1.46	1.38
4	D	315	UMP	O4'-C1'	4.93	1.53	1.42
4	D	314	UMP	O4'-C1'	5.07	1.54	1.42
3	C	414	D16	C-N	5.24	1.45	1.34
4	D	315	UMP	C6-C5	5.35	1.47	1.35
4	D	314	UMP	C6-C5	5.44	1.47	1.35
3	C	415	D16	C-N	5.76	1.46	1.34
3	C	414	D16	O4-C4	8.86	1.40	1.23
3	C	415	D16	O4-C4	9.62	1.42	1.23

All (52) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	415	D16	C15-C14-S13	-5.52	105.99	112.23
4	D	315	UMP	C4'-O4'-C1'	-5.42	95.50	109.42
4	D	314	UMP	C1'-N1-C6	-5.29	110.70	121.49
3	C	414	D16	C15-C14-S13	-5.24	106.31	112.23
4	D	315	UMP	C1'-N1-C6	-5.17	110.95	121.49
4	D	314	UMP	C4'-O4'-C1'	-5.12	96.27	109.42
4	D	314	UMP	O4-C4-C5	-4.69	117.83	125.28
4	D	315	UMP	O4-C4-C5	-4.58	118.00	125.28
3	C	415	D16	C15-C14-N10	-3.64	124.22	127.89
3	C	414	D16	C15-C14-N10	-3.12	124.74	127.89
4	D	314	UMP	C4-N3-C2	-3.02	122.25	126.52
3	C	414	D16	CP1-N10-C14	-2.95	116.13	120.74
3	C	415	D16	C16-C11-S13	-2.95	106.13	110.44
4	D	315	UMP	C4-N3-C2	-2.91	122.40	126.52
3	C	415	D16	CA-N-C	-2.76	117.38	122.01
4	D	314	UMP	O4'-C1'-C2'	-2.64	101.10	106.28
3	C	414	D16	OE2-CD-CG	-2.60	114.74	123.09
4	D	314	UMP	OP1-P-OP3	-2.58	102.20	110.63
4	D	315	UMP	OP1-P-OP3	-2.55	102.32	110.63
3	C	415	D16	OE2-CD-CG	-2.51	115.03	123.09
3	C	414	D16	O4-C4-N3	-2.50	118.18	120.42
3	C	414	D16	C4A-C8A-N1	-2.44	117.47	119.52
3	C	414	D16	CA-N-C	-2.40	117.99	122.01

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	314	UMP	C6-C5-C4	-2.37	116.55	119.54
3	C	415	D16	O4-C4-N3	-2.25	118.40	120.42
3	C	414	D16	C16-C11-S13	-2.14	107.32	110.44
3	C	415	D16	C4A-C8A-N1	-2.09	117.76	119.52
4	D	314	UMP	O2-C2-N3	-2.05	117.41	121.44
4	D	314	UMP	C2'-C3'-C4'	2.18	107.19	102.77
4	D	314	UMP	O5'-P-OP3	2.27	112.80	107.08
3	C	415	D16	CM2-C2-N1	2.28	121.04	116.53
4	D	314	UMP	OP2-P-O5'	2.38	113.68	106.72
4	D	315	UMP	N3-C2-N1	2.42	118.21	114.84
4	D	314	UMP	N3-C2-N1	2.42	118.21	114.84
3	C	414	D16	C4A-C4-N3	2.54	121.07	117.73
4	D	315	UMP	C2'-C3'-C4'	2.75	108.36	102.77
3	C	415	D16	OE1-CD-OE2	2.86	130.51	123.30
3	C	414	D16	OE1-CD-OE2	2.91	130.63	123.30
4	D	315	UMP	C5-C4-N3	3.28	119.83	114.86
4	D	314	UMP	C5-C4-N3	3.74	120.52	114.86
4	D	315	UMP	C2'-C1'-N1	4.71	124.99	113.52
4	D	314	UMP	O4'-C1'-N1	4.85	117.10	107.83
3	C	414	D16	C-C11-S13	5.18	126.83	115.77
4	D	315	UMP	C1'-N1-C2	5.27	129.16	117.63
4	D	314	UMP	C2'-C1'-N1	5.30	126.43	113.52
4	D	314	UMP	C1'-N1-C2	5.45	129.57	117.63
3	C	415	D16	C-C11-S13	5.92	128.41	115.77
4	D	315	UMP	O4'-C1'-N1	6.81	120.85	107.83
3	C	414	D16	C11-S13-C14	7.15	99.38	91.38
3	C	415	D16	C11-S13-C14	7.60	99.88	91.38
3	C	414	D16	S13-C14-N10	9.20	128.95	120.78
3	C	415	D16	S13-C14-N10	10.14	129.78	120.78

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	D	315	UMP	C1'
4	D	314	UMP	C1'

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	414	D16	3	0
3	C	415	D16	2	0
4	D	314	UMP	2	0
4	D	315	UMP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

PRELIMINARY VALIDATION REPORT

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9	
1	A	281/281 (100%)	-0.24	2 (0%)	89 88	39, 56, 71, 84	0
2	B	281/283 (99%)	-0.23	1 (0%)	93 93	39, 53, 68, 91	0
All	All	562/564 (99%)	-0.24	3 (0%)	91 91	39, 55, 71, 91	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	332	ALA	2.5
1	A	77	ILE	2.5
1	A	330	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	D16	C	415	32/?	0.89	0.18	1.94	46,61,89,94	0
3	D16	C	414	32/?	0.89	0.18	0.68	50,60,80,83	0
4	UMP	D	314	20/?	0.95	0.15	0.54	47,55,61,63	0
4	UMP	D	315	20/?	0.97	0.14	-0.04	47,52,57,59	0

6.5 Other polymers [i](#)

There are no such residues in this entry.

PRELIMINARY VALIDATION REPORT