	Sources of funding and other support (such as supply of drugs), role of funders	25	Funding
_	Where the full trial protocol can be accessed, if available	24	Protocol
	Registration number and name of trial registry	23	Registration
1	Interpretation consistent with results, balancing benefits and harms, and considering other relevant evidence	22	Interpretation
74	Generalisability (external validity, applicability) of the trial findings	2	Generalisability
13) 14	Trial limitations, addressing sources of potential bias, imprecision, and, if relevant, multiplicity of analyses	20	Discussion Limitations
13	All important harms or unintended effects in each group (for specific guidance see CONSORT for harms)	19	Harms
11, 12, 22, 8	pre-specified from exploratory		
	Results of any other analyses performed, including subgroup analyses and adjusted analyses, distinguishing	18	Ancillary analyses
11.12.22	For binary outcomes, presentation of both absolute and relative effect sizes is recommended	17b	
11,12	precision (such as 95% confidence interval)		estimation
	For each primary and secondary outcome, results for each group, and the estimated effect size and its	17a	Outcomes and
×	by original assigned groups		
3 -	For each group, number of participants (denominator) included in each analysis and whether the analysis was	16	Numbers analysed
9 20	A table showing baseline demographic and clinical characteristics for each group	15	Baseline data
5	Why the trial ended or was stopped	14b	
ý	Dates defining the periods of recruitment and follow-up	14a	Recruitment
~	For each group, losses and exclusions after randomisation, together with reasons	13b	recommended)
×	were analysed for the primary outcome		diagram is strongly
1	For each group, the numbers of participants who were randomly assigned, received intended treatment, and	13a	Participant flow (a
-			Results
9-12	Methods for additional analyses, such as subgroup analyses and adjusted analyses	12b	Statistical methods
0	If relevant, description of the similarity of interventions	11b	
7	assessing outcomes) and how		

recommend reading CONSORT extensions for cluster randomised trials, non-inferiority and equivalence trials, non-pharmacological treatments, herbal interventions, and pragmatic trials. Additional extensions are forthcoming: for those and for up to date references relevant to this checklist, see www.consort-statement.org. *We strongly recommend reading this statement in conjunction with the CONSORT 2010 Explanation and Elaboration for important clarifications on all the items. If relevant, we also