

S2 Table

	SHAM	NX	NX LIRA
N	14	14	15
Lira-treatment baseline			
Body weight (g)	20.4 ± 0.3	20.3 ± 0.5	19.8 ± 0.3
P-urea (mmol/L)	10.3 ± 0.3	30.5 ± 2.6 ^a	27.6 ± 2.3
P-cholesterol (mmol/L)	6.5 ± 0.2	7.7 ± 0.2 ^b	7.3 ± 0.2
P-phosphate	1.9 ± 0.08	1.8 ± 0.05	1.8 ± 0.08
Termination			
Body weight (g)	21.7 ± 0.3	21.2 ± 0.2	19.8 ± 0.3 ^c
P-urea (mmol/L)	9.4 ± 0.4	20.5 ± 1.4 ^a	19.9 ± 1.6
P-creatinine (μmol/L)	15.3 ± 1.3	24.1 ± 1.2 ^a	20.6 ± 1.2
P-cholesterol (mmol/L)	40.2 ± 1.3	40.7 ± 1.6	41.9 ± 2.7
P-phosphate	1.8 ± 0.09	1.9 ± 0.09	1.9 ± 0.11
P- Ca ²⁺ (mmol/L)	2.2 ± 0.02	2.5 ± 0.03 ^a	2.5 ± 0.04

Results are depicted as mean±SEM. Statistical analysis was determined by 1-way ANOVA followed by Sidak's multiple comparisons post-test.

^a=**** (SHAM vs NX)

^b=*** (SHAM vs NX)

^c=** (NX vs NX LIRA)